OPDIVO[®] (nivolumab) OPDIVO + YERVOY[®] (nivolumab) (ipilimumab)

Patient Alert Card

This material was developed by Bristol-Myers Squibb as part of the risk minimization plan for OPDIVO. This material is not intended for promotional use.

Date of Internal Approval: September 2020

Local Approval Number: 7356CA2000183

\mu Bristol Myers Squibb

Important Information for Patients

Please carry this card with you at all times to inform healthcare professionals that you are receiving treatment with OPDIVO or OPDIVO in combination with YERVOY.

i) IMPORTANT

- Tell your doctor of any previous medical conditions, including if you have had a stem cell transplant that uses donor stem cells (allogeneic).
- Early assessment and management of side effects by your doctor is critical.
- Signs and symptoms that may appear mild can quickly worsen if left untreated.
- **DO NOT** try to treat these symptoms yourself.
- Signs and symptoms may be delayed and may occur weeks to months after your last injection.

For more information, consult the OPDIVO Product Monograph https://www.bms.com/assets/bms/ ca/documents/productinformation/ OPDIVO_EN_PIL.pdf and the Patient Information Guide or call Medical Information at 1-866-463-6267.


D POSSIBLE SIDE EFFECTS

OPDIVO treatment may increase the risk of serious or even life-threatening immune-mediated side effects, which may affect different parts of the body. Signs and symptoms to look out for include the following, though these are not the only ones:



GENERAL:

- Changes in mood or behaviour (e.g., decreased sex drive, irritability, forgetfulness, depression),
- confusion and memory problems,
- feeling unwell,
- fever,
- headaches,
- increased or decreased appetite,
- nausea or vomiting,
- seizures (fits),
- seeing things that are not really there (hallucination),
- swelling in extremities,
- swollen lymph nodes,
- excessive thirst,
- hunger,
- sensation of paralysis,
- pain on the right side of your stomach (abdomen),
- tiredness,
- weight gain or loss,
- tingling and/or numbness (e.g., in your fingers or toes),
- sleepiness or drowsiness



MUSCLES AND BONES:

- Difficulty walking,
- numbness in arms and/or legs,
- muscle or joint pains,
- muscle weakness or stiffness,
- stiff neck

KIDNEYS AND BLADDER:

- Decreased or increased amount of urine,
- darker urine than normal (tea-coloured)

STOMACH AND BOWEL (GUT):


- Dark, tarry, or sticky stools,
- blood or mucous in your stools,
- diarrhea (watery, loose or soft stools) or more bowel movements than usual,
- heartburn or indigestion,
- pain or tenderness in your stomach or abdominal area



- Eyesight changes (e.g., blurry vision, other vision problems),
- eye pain or redness,
- yellowing of the whites of your eyes



- Chest pain,
- breathing difficulties,
- cough,
- irregular heartbeat,
- palpitations (being more aware of heartbeat than normal)



• Ulcers in the mouth or other mucous membranes



- Bruising easily,
- dry skin,
- itching,
- raised skin lumps/bumps (skin nodules),
- severe skin reaction or rash,
- skin blistering/peeling,
- skin yellowing (jaundice)



- This patient is treated with OPDIVO or OPDIVO in combination with YERVOY.
- Immune-mediated adverse reactions (imARs) may appear at any time during treatment or months after its discontinuation.
- Early diagnosis and appropriate management are essential to minimize life-threatening complications.

Reporting side effects

You can report any suspected side effects associated with the use of Health Products to Health Canada by:

 Visiting the Web page on Adverse Reaction Reporting (Adverse Reaction Reporting (https://www.canada.ca/en/ health-canada/services/ drugs-health-products/ medeffect-canada/adversereaction-reporting.html) for information on how to report online, by mail or by fax; or • Consultation with an oncologist or other medical specialist may be helpful for management of organ-specific imARs.

Consult the OPDIVO Product Monograph at https://www.bms.com/assets/ bms/ca/documents/ productmonograph/OPDIVO_ EN_PM.pdf

or call Medical Information at 1-866-463-6267 for more information.

 Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice. My Doctor's Contact Information (who prescribed OPDIVO or OPDIVO in combination with YERVOY)

Name of Doctor:

Office Phone:

After-Hours Phone:

The healthcare professional treating this patient with OPDIVO or OPDIVO in combination with YERVOY should complete the 'My doctor's contact information' section of this Patient Alert Card.

My Contact Information

My Name and Phone Number:

Emergency Contact (name and phone):

OPDIVO and YERVOY are registered trademarks of Bristol-Myers Squibb Company used under license by Bristol-Myers Squibb Canada Co.

©2020 Bristol-Myers Squibb Company. All rights reserved.

OPDIVO and YERVOY are registered trademarks of Bristol-Myers Squibb Company used under license by Bristol-Myers Squibb Canada Co.

