

## PRODUCT MONOGRAPH

Pr **SUSTIVA**<sup>®</sup>

(efavirenz, USP)

efavirenz Capsules USP, 50 and 200 mg

Antiretroviral Agent

Bristol-Myers Squibb Canada  
Montreal, Canada

Date of Preparation:  
March 18, 1999

<sup>®</sup> Registered trademark of Bristol-Myers  
Squibb Pharma Company used under licence by  
Bristol-Myers Squibb Canada

Date of Revision:  
August 16, 2023

Control Number: 273530

## TABLE OF CONTENTS

<b>PRODUCT MONOGRAPH .....</b>	<b>1</b>
TABLE OF CONTENTS .....	2
<b>PART I: HEALTH PROFESSIONAL INFORMATION .....</b>	<b>3</b>
SUMMARY PRODUCT INFORMATION.....	3
INDICATIONS AND CLINICAL USE .....	3
CONTRAINDICATIONS.....	3
WARNINGS AND PRECAUTIONS .....	4
ADVERSE REACTIONS .....	11
DRUG INTERACTIONS .....	17
DOSAGE AND ADMINISTRATION .....	23
OVERDOSAGE.....	24
ACTION AND CLINICAL PHARMACOLOGY.....	25
STORAGE AND STABILITY .....	32
DOSAGE FORMS, COMPOSITION AND PACKAGING.....	33
<b>PART II: SCIENTIFIC INFORMATION.....</b>	<b>34</b>
PHARMACEUTICAL INFORMATION .....	34
CLINICAL TRIALS .....	35
DETAILED PHARMACOLOGY .....	41
VIROLOGY .....	43
TOXICOLOGY .....	44
<b>PART III: CONSUMER INFORMATION.....</b>	<b>51</b>

Pr SUSTIVA®  
(efavirenz, USP)

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form / Strength</b>	<b>Clinically Relevant Nonmedicinal Ingredients*</b>
Oral	Capsules / 50 and 200 mg	lactose monohydrate

\* For a complete listing, see [DOSAGE FORMS, COMPOSITION AND PACKAGING](#) section

**INDICATIONS AND CLINICAL USE**

SUSTIVA (efavirenz) is indicated for the treatment of HIV-1 infection in combination with other antiretroviral agents.

**CONTRAINDICATIONS**

SUSTIVA (efavirenz) is contraindicated in patients with clinically significant hypersensitivity to any of its components. For a complete listing, see the [DOSAGE FORMS, COMPOSITION AND PACKAGING](#) section of the Product Monograph.

SUSTIVA must not be administered concurrently with cisapride<sup>1</sup>, midazolam, triazolam, pimozide or ergot derivatives because competition for CYP3A4 by efavirenz could result in inhibition of metabolism of these drugs and create the potential for serious and/or life-threatening adverse events (e.g., cardiac arrhythmias, prolonged sedation or respiratory depression). (See [Table 1](#)).

Coadministration of efavirenz with elbasvir and grazoprevir is contraindicated, due to the potential for significant decreases in plasma concentrations of elbasvir and grazoprevir. This effect is due to an induction of CYP3A4 by efavirenz, and may result in loss of therapeutic effect

Herbal preparations containing St. John's wort (*Hypericum perforatum*) must not be used while taking SUSTIVA due to the risk of decreased plasma concentrations and reduced clinical effects of SUSTIVA.

---

<sup>1</sup> Cisapride is not marketed in Canada

**Table 1**

<b>Drugs That Are Contraindicated with SUSTIVA</b>		
<b>Drug Class</b>	<b>Drugs Within Class Contraindicated With SUSTIVA</b>	<b>Clinical Comment</b>
Benzodiazepines	midazolam, triazolam	Potential for serious and/or life-threatening reactions such as prolonged or increased sedation or respiratory depression.
GI Motility Agents	cisapride*	Potential for serious and/or life-threatening reactions such as cardiac arrhythmias.
Anti-Migraine	ergot derivatives (dihydroergotamine, ergonovine, ergotamine)	Potential for serious and/or life-threatening reactions such as acute ergot toxicity characterized by peripheral vasospasm and ischemia of the extremities and other tissues.
Neuroleptic	pimozide	Potential for serious and/or life-threatening reactions such as cardiac arrhythmias.
Hepatitis C antiviral agents	elbasvir/grazoprevir	May lead to loss of therapeutic effect of elbasvir/grazoprevir.
Herbal Products	St. John's wort ( <i>Hypericum perforatum</i> )	May lead to loss of virologic response and possible resistance to SUSTIVA or to the class of non-nucleoside reverse transcriptase inhibitors (NNRTIs).

\*not marketed in Canada

## WARNINGS AND PRECAUTIONS

### General

Coadministration of SUSTIVA with ATRIPLA, a fixed-dose combination tablet containing efavirenz, emtricitabine and tenofovir disoproxil fumarate, is not recommended, unless needed for dose adjustment (e.g, with rifampin).

### Effects on Ability to Drive and To Use Machines:

SUSTIVA may cause dizziness, impaired concentration, and/or drowsiness. Patients should be instructed that, if they experience these symptoms, they should avoid potentially hazardous tasks such as driving or operating machinery (see [Neurologic](#)).

### Carcinogenesis, Mutagenesis and Impairment of Fertility (see Part II: [TOXICOLOGY](#))

### Cardiovascular

QTc prolongation has been observed with the use of efavirenz (see [DRUG INTERACTIONS](#) and [ACTION AND CLINICAL PHARMACOLOGY](#)). Consider alternatives to SUSTIVA when coadministered with a drug with a known risk of Torsade de Pointes or when administered to patients at higher risk of Torsade de Pointes.

## Endocrine and Metabolism

### Serum Lipids and Blood Glucose:

Serum lipid and blood glucose levels may increase during antiretroviral therapy (ART). Disease control and lifestyle changes may also be contributing factors. Consideration should be given to the measurement of serum lipids and blood glucose. Lipid disorders and blood glucose elevations should be managed as clinically appropriate.

### Hepatic/Biliary/Pancreatic

Postmarketing reports of hepatic failure have occurred including some cases in patients without pre-existing hepatic disease or other identifiable risk factors (see [Monitoring and Laboratory Tests](#); [Liver Enzymes](#)).

In controlled clinical studies the rate of clinical pancreatitis was similar in patients receiving 1/1008 (0.1%) and not receiving efavirenz 2/635 (0.3%).

Asymptomatic increases in serum amylase levels were observed in a significantly higher number of patients treated with efavirenz 600 mg than in control patients (see [ADVERSE REACTIONS; Abnormal Hematologic and Clinical Chemistry Findings](#)).

Elevated triglycerides have been reported in patients receiving efavirenz, in some cases to levels which can predispose a patient to pancreatitis. Among patients with elevated triglycerides, there have been no cases of pancreatitis. Because these triglyceride levels were not obtained in a fasting state, the exact clinical relevance of these measurements is not known.

### Hypersensitivity Reactions

In clinical trials, hypersensitivity reactions were uncommon (<1%) in patients treated with SUSTIVA.

### Immune

Immune Reconstitution Inflammatory Syndrome has been reported in patients treated with combination antiretroviral therapy, including SUSTIVA. During the initial phase of treatment, patients responding to antiretroviral therapy may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium* infection, cytomegalovirus, *Pneumocystis jirovicii* pneumonia (PCP), or tuberculosis), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis and Guillain-Barre syndrome and autoimmune hepatitis) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable, and can occur many months after initiation of treatment.

## Neurologic

### Nervous System Symptoms:

Fifty-three percent of patients receiving SUSTIVA in controlled clinical trials reported central nervous system symptoms compared to 25% of patients receiving control regimens. These symptoms included, but were not limited to, dizziness (28.1%), insomnia (16.3%), impaired concentration (8.3%), somnolence (7.0%), abnormal dreams (6.2%), and hallucinations (1.2%). In controlled trials, these symptoms were severe in 2.0% of patients receiving SUSTIVA 600 mg daily and in 1.3% of patients receiving control regimens. In clinical trials, 2.1% of SUSTIVA - treated patients discontinued therapy because of nervous system symptoms. These symptoms usually begin during the first or second day of therapy and generally resolve after the first 2-4 weeks. After 4 weeks of therapy the prevalence of nervous system symptoms of at least moderate severity ranged from 5-9% in patients treated with regimens containing SUSTIVA and from 3-5% in patients treated with a control regimen. Patients should be informed that these common nervous system symptoms are likely to improve with continued therapy. Dosing at bedtime improves tolerability of these symptoms (see [ADVERSE REACTIONS](#) and [DOSAGE AND ADMINISTRATION](#)).

Analysis of long-term data from Study AI266-006 (median follow-up 180 weeks, 102 weeks, and 76 weeks for patients treated with SUSTIVA + zidovudine + lamivudine, SUSTIVA + indinavir, and indinavir + zidovudine + lamivudine, respectively) showed that, beyond 24 weeks of therapy, the incidences of new-onset nervous system symptoms among SUSTIVA-treated patients were generally similar to those in the indinavir-containing control arm.

Patients receiving SUSTIVA should be alerted to the potential for additive central nervous system effects when SUSTIVA is used concomitantly with alcohol or psychoactive drugs.

### Seizures:

Caution should be taken in any patient with a history of seizures. Convulsions have been observed infrequently in patients receiving efavirenz, generally in the presence of known medical history of seizures. Overall, the rate of seizure in controlled clinical trials has been 0.89% in SUSTIVA treated patients and 0.63% in the control patients. Patients who are receiving concomitant anticonvulsant medications primarily metabolized by the liver such as phenytoin, carbamazepine, and phenobarbital, may require periodic monitoring of plasma levels (see [DRUG INTERACTIONS](#)).

### Psychiatric:

Serious psychiatric adverse experiences have been reported in patients treated with SUSTIVA. In controlled trials of 1008 patients treated with regimens containing SUSTIVA for an average of 2.1 years and 635 patients treated with control regimens for an average of 1.5 years, the frequency of specific serious psychiatric events among patients who received efavirenz or control regimens respectively, were: severe depression (2.4%, 0.9%), suicidal ideation (0.7%, 0.3%), non-fatal suicide attempts (0.5%, 0%), aggressive behavior (0.4%, 0.5%), paranoid

reactions (0.4%, 0.3%), and manic reactions (0.2%, 0.3%). When psychiatric symptoms similar to those noted above were combined and evaluated as a group in a multifactorial analysis of data from Study AI266-006, treatment with efavirenz was associated with an increase in the occurrence of these selected psychiatric symptoms. Other factors associated with an increase in the occurrence of these psychiatric symptoms were history of injection drug use, psychiatric history, and receipt of psychiatric medication at study entry; similar associations were observed in both the SUSTIVA and control treatment groups. In Study AI266-006, onset of new serious psychiatric symptoms occurred throughout the study for both SUSTIVA-treated and control-treated patients. One percent of SUSTIVA-treated patients discontinued or interrupted treatment because of one or more of these selected psychiatric symptoms. There have also been occasional post marketing reports of death by suicide, delusions, psychosis – like behavior and catatonia, although a causal relationship to the use of SUSTIVA cannot be determined from these reports. Patients with serious psychiatric adverse experiences should seek immediate medical evaluation to assess the probability that the symptoms may be related to the use of SUSTIVA, and if so, to determine whether the risks of continued therapy outweigh the benefits (see [ADVERSE REACTIONS](#)).

#### **Sensitivity/Resistance:**

SUSTIVA (efavirenz) must not be used as a single agent to treat HIV or added on as a sole agent to a failing regimen. As with all other non-nucleoside reverse transcriptase inhibitors, resistant virus emerges rapidly when efavirenz is administered as monotherapy. The choice of new antiretroviral agents to be used in combination with efavirenz should take into consideration the potential for viral cross-resistance. (Please refer to the most recent antiretroviral guidelines for further information.)

#### **Resistance:**

Clinical isolates with reduced susceptibility in cell culture to efavirenz have been obtained. The most frequently observed amino acid substitution in clinical studies with efavirenz is K103N (54%). One or more RT substitutions at amino acid positions 98, 100, 101, 103, 106, 108, 188, 190, 225, 227, and 230 were observed in patients failing treatment with efavirenz in combination with other antiretrovirals. Other resistance mutations observed to emerge commonly included L100I (7%), K101E/Q/R (14%), V108I (11%), G190S/T/A (7%), P225H (18%), and M230I/L (11%) (see [VIROLOGY](#)).

#### **Cross-Resistance:**

Cross-resistance has been recognized among NNRTIs. Clinical isolates previously characterized as efavirenz-resistant were also phenotypically resistant in cell culture to delavirdine and nevirapine compared to baseline. Delavirdine- and/or nevirapine-resistant clinical viral isolates with NNRTI resistance-associated substitutions (A98G, L100I, K101E/P, K103N/S, V106A, Y181X, Y188X, G190X, P225H, F227L, or M230L) showed reduced susceptibility to efavirenz in cell culture (see [VIROLOGY](#)).

**Skin:**

Efavirenz is not recommended for patients who have had a life-threatening cutaneous reaction (eg, Stevens-Johnson syndrome). SUSTIVA should be discontinued in patients developing severe rash associated with blistering, desquamation, mucosal involvement or fever.

In controlled clinical trials, 26% (266/1008) of patients treated with 600 mg SUSTIVA experienced new onset skin rash compared with 17% (111/635) of patients treated in control groups. Rash associated with blistering, moist desquamation or ulceration occurred in 0.9% (9/1008) of patients treated with SUSTIVA. The median time to onset of rash in adults was 11 days and the median duration, 16 days. The discontinuation rate for rash in clinical trials was 6.4% (17/266) among patients with rash and 1.7% (17/1008) overall.

In clinical trials, grade 4 rash (including Stevens-Johnson syndrome, erythema multiforme, toxic epidermal necrolysis and exfoliative dermatitis) was uncommon (<1%) in patients treated with SUSTIVA.

Rash was reported in 26 of 57 pediatric patients (46%) treated with SUSTIVA capsules. One pediatric patient experienced Grade 3 rash (confluent rash with fever), and two patients had Grade 4 rash (erythema multiforme). The median time to onset of rash in children was eight days. Prophylaxis with appropriate antihistamines prior to initiating therapy with SUSTIVA in children may be considered (see [ADVERSE REACTIONS](#)).

**Special Populations****Pregnant Women:**

Efavirenz may cause fetal harm when administered during the first trimester to a pregnant woman. Pregnancy should be avoided in women receiving SUSTIVA and for 12 weeks after discontinuation. Barrier contraception must always be used in combination with other methods of contraception (e.g., oral or other hormonal contraceptives) (see [DRUG INTERACTIONS](#) and [TOXICOLOGY](#)). Because of the long half-life of efavirenz, use of adequate contraceptive measures for 12 weeks after discontinuation of SUSTIVA is recommended. Women of childbearing potential should undergo pregnancy testing prior to initiation of SUSTIVA (see Reproductive Risk Potential).

There are no adequate and well-controlled studies in pregnant women. SUSTIVA should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus, such as in pregnant women without other therapeutic options.

**Antiretroviral Pregnancy Registry**

To monitor fetal outcomes of pregnant women exposed to SUSTIVA, an Antiretroviral Pregnancy Registry has been established. Physicians are encouraged to register patients,

<http://www.apregistry.com>

Telephone: (800) 258-4263

Fax: (800) 800-1052

As of July 2013, the Antiretroviral Pregnancy Registry has received prospective reports of 1067 pregnancies exposed to efavirenz-containing regimens, 904 of which were first-trimester exposures. Birth defects occurred in 18 of 766 live births (first-trimester exposure) and 3 of 160 live births (second-/third-trimester exposure). One of these prospectively reported defects with first-trimester exposure was a neural tube defect. A single case of anophthalmia with first trimester exposure to efavirenz has also been prospectively reported; however, this case included severe oblique facial clefts and amniotic banding, a known association with anophthalmia.

There have been seven reports during post-marketing use of findings consistent with neural tube defects, including meningocele, all in mothers exposed to efavirenz containing regimens (excluding any efavirenz-containing fixed-dose combination tablets) in the first trimester. Although a causal relationship of these events to the use of SUSTIVA cannot be established, similar defects have been observed in nonclinical studies of efavirenz (see [TOXICOLOGY, Reproduction and Teratology](#)).

#### **Nursing Women:**

It is recommended that women living with HIV do not breast-feed their infants in order to avoid transmission of HIV. Efavirenz has been shown to be excreted in human milk. There is insufficient information on the effects of efavirenz in newborns/infants. Risk to the infant cannot be excluded. Breast-feeding should be discontinued during treatment with SUSTIVA.

#### **Pediatrics:**

ACTG 382 is an ongoing open-label uncontrolled 48-week study in 57 NRTI-experienced pediatric patients to characterize the safety, pharmacokinetics, and antiviral activity of SUSTIVA in combination with nelfinavir (20-30 mg/kg TID) and NRTIs. Mean age was 8 years (range 3-16). SUSTIVA has not been studied in pediatric patients below 3 years of age or who weigh less than 13 kg. At 48 weeks, the type and frequency of adverse experiences was generally similar to that of adult patients with the exception of a higher incidence of rash which was reported in 46% (26/57) of pediatric patients compared to 26% of adults, and a higher frequency of Grade 3 or 4 rash reported in 5% (3/57) of pediatric patients compared to 0.9% of adults (see [ADVERSE REACTIONS](#)).

The starting dose of SUSTIVA was 600 mg daily adjusted to body size, based on weight, targeting AUC levels in the range of 190-380  $\mu\text{M}\cdot\text{h}$ . The pharmacokinetics of efavirenz in pediatric patients were similar to adults. In 48 pediatric patients receiving the equivalent of a 600 mg dose of SUSTIVA, steady-state  $C_{\text{max}}$  was  $14.2 \pm 5.8 \mu\text{M}$  (mean  $\pm$  SD), steady-state  $C_{\text{min}}$  was  $5.6 \pm 4.1 \mu\text{M}$ , and AUC was  $218 \pm 104 \mu\text{M}\cdot\text{h}$  (see also [DETAILED PHARMACOLOGY](#)).

#### **Geriatrics:**

Clinical studies of SUSTIVA did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. In general, dose selection for

an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other therapy.

### **Hepatic Impairment:**

The pharmacokinetics of efavirenz have not been adequately studied in patients with hepatic impairment and its use is not recommended in patients with moderate or severe hepatic impairment (Child-Pugh Class B or C) because of insufficient data. Because of the extensive cytochrome P450-mediated metabolism of efavirenz and limited clinical experience in patients with mild hepatic impairment, caution should be exercised in administering SUSTIVA to these patients (see [Monitoring and Laboratory Tests](#); [Liver Enzymes](#), [ADVERSE REACTIONS](#); [Abnormal Hematologic and Clinical Chemistry Findings](#), [DOSAGE AND ADMINISTRATION](#) and [ACTION AND CLINICAL PHARMACOLOGY](#); [Pharmacokinetics](#); [Special Populations](#)).

### **Renal Impairment:**

The pharmacokinetics of efavirenz have not been studied in patients with renal insufficiency. However, less than 1% of efavirenz is excreted unchanged in the urine; consequently, the impact of renal impairment on efavirenz elimination should be minimal. There is no experience in patients with severe renal failure and close safety monitoring is recommended in this population (see [DOSAGE AND ADMINISTRATION](#)).

### **Monitoring and Laboratory Tests**

#### **Lipids:**

Monitoring of cholesterol and triglycerides should be considered in patients treated with SUSTIVA (see [ADVERSE REACTIONS](#); [Abnormal Hematologic and Clinical Chemistry Findings](#)).

#### **Liver Enzymes:**

In patients with underlying liver disease including Hepatitis B or C infection and in patients treated with other medications associated with liver toxicity, monitoring of liver enzymes is recommended. Postmarketing reports of hepatic failure have occurred including some cases in patients with no pre-existing hepatic disease or other identifiable risk factors (see [Post-Market Adverse Drug Reactions](#)). Liver enzyme monitoring should also be considered for patients without pre-existing hepatic dysfunction or other risk factors. In patients with persistent elevations of serum transaminases to greater than five times the upper limit of the normal range, the benefit of continued therapy with SUSTIVA needs to be weighed against the potential risks of significant liver toxicity (see [ADVERSE REACTIONS](#); [Abnormal Hematologic and Clinical Chemistry Findings](#)).

## ADVERSE REACTIONS

### Adverse Drug Reaction Overview

SUSTIVA (efavirenz) has been studied in 9200 patients. The most significant adverse events observed in patients treated with SUSTIVA are nervous system symptoms, psychiatric symptoms, and rash.

The long-term safety profile of SUSTIVA-containing regimens was evaluated in a controlled trial, in which patients received SUSTIVA + zidovudine + lamivudine (n = 412, median duration 180 weeks), SUSTIVA + indinavir (n = 415, median duration 102 weeks), or indinavir + zidovudine + lamivudine (n = 401, median duration 76 weeks). Long-term use of SUSTIVA in this study was not associated with any new safety concerns.

### Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse drug reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

#### Nervous System Symptoms

Fifty-three percent of patients receiving SUSTIVA reported central nervous system symptoms (see [WARNINGS AND PRECAUTIONS; Neurologic](#)). Table 2 lists the frequency of the symptoms of different degrees of severity, and gives the discontinuation rates in clinical trials for one or more of the following nervous system symptoms: dizziness, insomnia, impaired concentration, somnolence, abnormal dreaming, euphoria, confusion, agitation, amnesia, hallucinations, stupor, abnormal thinking, and depersonalization. The frequencies of specific central and peripheral nervous system symptoms are provided in Table 2.

**Table 2**  
**Percent of Patients with One or More Selected Nervous System Symptoms<sup>1,2</sup>**

Percent of Patients with:	SUSTIVA 600 mg daily (N=1008)	Control Groups (N= 635)
	%	%
Mild Symptoms <sup>3</sup>	33.3	15.6
Moderate Symptoms <sup>4</sup>	17.4	7.7
Severe Symptoms <sup>5</sup>	2	1.3
Symptoms of Any Severity	52.7	24.6
Treatment discontinuation as a result of symptoms	2.1	1.1

<sup>1</sup> Includes events reported regardless of causality.

<sup>2</sup> Data from Studies 006, 020 and two Phase II studies.

<sup>3</sup> "Mild" = Symptoms which do not interfere with patient's daily activities.

<sup>4</sup> "Moderate" = Symptoms which may interfere with daily activities.

<sup>5</sup> "Severe" = Symptoms which interrupt patient's usual daily activities.

Analysis of long-term data (median treatment duration 180 weeks, 102 weeks, and 76 weeks for patients treated with SUSTIVA + zidovudine + lamivudine, SUSTIVA + indinavir, and indinavir + zidovudine + lamivudine, respectively) showed that, beyond 24 weeks of therapy, the incidences of new-onset nervous system symptoms among SUSTIVA-treated patients were generally similar to those in the control arm.

### Psychiatric Symptoms

Serious psychiatric adverse experiences have been reported in patients treated with SUSTIVA. In controlled trials the frequency of specific serious psychiatric symptoms among patients who received SUSTIVA or control regimens, respectively, were: severe depression (2.4%, 0.9%), suicidal ideation (0.7%, 0.3%), non-fatal suicide attempts (0.5%, 0%), aggressive behavior (0.4%, 0.5%), paranoid reactions (0.4%, 0.3%) and manic reactions (0.2%, 0.3%) (see [WARNINGS AND PRECAUTIONS; Psychiatric](#)). Additional psychiatric symptoms observed at a frequency of >2% among patients treated with SUSTIVA or control regimens respectively, in controlled clinical trials were depression (19%, 16%), anxiety (13%, 9%) and nervousness (7%, 2%).

### Skin Rash

Rashes are usually mild-to-moderate maculopapular skin eruptions that occur within the first two weeks of initiating therapy with SUSTIVA. In most patients rash resolves with continuing SUSTIVA therapy within one month. SUSTIVA can be reinitiated in patients interrupting therapy because of Grades 1 and 2 rash. SUSTIVA should be discontinued in patients developing severe rash associated with blistering, desquamation, mucosal involvement or fever. The frequency of rash by NCI grade and the discontinuation rates as a result of rash are provided in Table 3.

**Table 3**  
**Percent of Patients with Treatment-Emergent Rash<sup>1,2</sup>**

<b>Percent of Patients with:</b>	<b>Description of Rash Grade3</b>	<b>SUSTIVA 600 mg once daily Adults (N= 1008) %</b>	<b>SUSTIVA Pediatric Patients (N=57) %</b>	<b>Control Groups Adults (N= 635) %</b>
Grade 1 Rash	Erythema, pruritus	10.7	8.8	9.8
Grade 2 Rash	Diffuse maculopapular rash, dry desquamation	14.7	31.6	7.4
Grade 3 Rash	Vesiculation, moist desquamation, ulceration	0.8	1.8	0.3

Percent of Patients with:	Description of Rash Grade <sup>3</sup>	SUSTIVA 600 mg once daily Adults (N= 1008) %	SUSTIVA Pediatric Patients (N=57) %	Control Groups Adults (N= 635) %
Grade 4 Rash	Erythema multiforme, Stevens-Johnson Syndrome, toxic epidermal necrolysis, necrosis requiring surgery, exfoliative dermatitis	0.1	3.5	0
Rash of Any Grade	—	26.3	45.6	17.5
Treatment discontinuation as a result of rash	—	1.7	8.8	0.3

<sup>1</sup> Includes events reported regardless of causality.

<sup>2</sup> Data from Studies 006, 020, and two Phase II studies.

<sup>3</sup> NCI (National Cancer Institute) Grading System.

As seen in Table 3, rash is more common in children and more often of higher grade (i.e., more severe) (see [WARNINGS AND PRECAUTIONS; Skin](#)).

Experience with SUSTIVA in patients who discontinued other antiretroviral agents of the NNRTI class is limited. Nineteen patients who discontinued nevirapine because of rash have been treated with SUSTIVA. Nine of these patients developed mild-to-moderate rash while receiving therapy with SUSTIVA, and two of these patients discontinued because of rash.

Selected clinical adverse experiences of moderate or severe intensity observed in  $\geq 2\%$  of SUSTIVA-treated patients in three controlled clinical trials are presented in Table 4.

**Table 4**  
**Selected Treatment-Emergent<sup>1</sup> Adverse Events of**  
**Moderate or Severe Intensity Reported in  $\geq 2\%$  of SUSTIVA-Treated Patients**  
**in Studies DMP 266-006, ACTG 364 and DMP 266-020**

Adverse Events	Study DMP 266-006 3TC, NNRTI and Protease Inhibitor- Naive Patients			Study ACTG 364 NRTI-experienced NNRTI and Protease Inhibitor-Naive Patients			Study DMP 266-020 NRTI-experienced NNRTI and Protease Inhibitor-Naive Patients	
	SUSTIVA <sup>2</sup> + ZDV/3TC (N = 412) 180 weeks <sup>3</sup>	SUSTIVA <sup>2</sup> + Indinavir (N = 415) 102 weeks <sup>3</sup>	Indinavir + ZDV/3TC (N = 401) 76 weeks <sup>3</sup>	SUSTIVA <sup>2</sup> + Nelfinavir + NRTIs (N = 64)	SUSTIVA <sup>2</sup> + NRTIs (N = 65)	Nelfinavir + NRTIs (N = 66)	SUSTIVA <sup>2</sup> + Indinavir + NRTIs (N = 154)	Indinavir + NRTIs (N = 168)
	%	%	%	%	%	%	%	%
<b>Body as a Whole</b>								
Fatigue	8	5	9	0	2	3	5	1
Pain	1	2	8	13	6	17	4	3
<b>Central and Peripheral Nervous System</b>								
Dizziness	9	9	2	2	6	6	7	1
Headache	8	5	3	5	2	3	5	4
<b>Gastrointestinal</b>								
Nausea	10	6	24	3	2	2	10	10
Vomiting	6	3	14	-	-	-	6	5
Diarrhea	3	5	6	14	3	9	11	3

Adverse Events	Study DMP 266-006 3TC, NNRTI and Protease Inhibitor- Naive Patients			Study ACTG 364 NRTI-experienced NNRTI and Protease Inhibitor-Naive Patients			Study DMP 266-020 NRTI-experienced NNRTI and Protease Inhibitor-Naive Patients	
	SUSTIVA <sup>2</sup> + ZDV/3TC (N = 412) 180 weeks <sup>3</sup>	SUSTIVA <sup>2</sup> + Indinavir (N = 415) 102 weeks <sup>3</sup>	Indinavir + ZDV/3TC (N = 401) 76 weeks <sup>3</sup>	SUSTIVA <sup>2</sup> + Nelfinavir + NRTIs (N = 64)	SUSTIVA <sup>2</sup> + NRTIs (N = 65)	Nelfinavir + NRTIs (N = 66)	SUSTIVA <sup>2</sup> + Indinavir + NRTIs (N = 154)	Indinavir + NRTIs (N = 168)
	%	%	%	%	%	%	%	%
Dyspepsia	4	4	6	0	0	2	3	1
Abdominal Pain	2	2	5	3	3	3	3	1
<b>Psychiatric</b>								
Concentration Impaired	5	3	<1	0	0	0	3	1
Insomnia	7	7	2	0	0	2	3	1
Anxiety	2	4	<1	-	-	-	2	1
Abnormal Dreams	3	1	0	-	-	-	2	1
Somnolence	2	2	<1	0	0	0	2	2
Depression	5	4	<1	3	0	5	2	0
Anorexia	1	<1	<1	0	2	2	5	1
Nervousness	2	2	0	2	0	2	1	0
<b>Skin &amp; Appendages</b>								
Rash	11	16	5	9	5	9	10	6
Pruritus	<1	1	1	9	5	9	2	1

<sup>1</sup> Includes those adverse events at least possibly related to study drug or of unknown relationship for Studies 006 and 020. Includes all adverse events regardless of relationship to study drug for Study ACTG 364.

<sup>2</sup> SUSTIVA provided as 600 mg once daily.

<sup>3</sup> Median duration of treatment

- Not Specified

ZDV = zidovudine, 3TC = lamivudine

Lipodystrophy (any severity, regardless of relationship to study regimen) was reported in 3%, 4%, and 5% of patients treated with SUSTIVA + zidovudine + lamivudine, SUSTIVA + indinavir, and indinavir + zidovudine + lamivudine, respectively. The frequencies of other adverse event terms that may be associated with lipodystrophy (abdomen enlarged, breast enlargement, cachexia, gynecomastia, lipodosis, lipoma, and obesity) ranged from <1% to 3% and were similar among the treatment groups.

Clinical adverse experiences observed in  $\geq 10\%$  of 57 pediatric patients aged 3 to 16 years who received SUSTIVA capsules, nelfinavir, and one or more NRTIs were: rash (46%), diarrhea/loose stools (39%), fever (21%), cough (16%), dizziness/lightheaded/fainting (16%), ache/pain/discomfort (14%), nausea/vomiting (12%), and headache (11%). The incidence of nervous system symptoms was 18% (10/57). One patient experienced Grade 3 rash, two patients had Grade 4 rash, and five patients (9%) discontinued because of rash (see [WARNINGS AND PRECAUTIONS](#); [Special Populations](#), [Pediatrics](#)).

Adverse clinical experiences of moderate to severe intensity observed in less than 2% of patients receiving SUSTIVA in all Phase II/III studies, including the North American expanded access program as well as post-marketing spontaneous reports, and considered at least possibly related or of unknown relationship to treatment are listed below by body system:

Body as a Whole: alcohol intolerance, allergic reaction, asthenia, fever, hot flushes, influenza-like symptoms, malaise, pain, peripheral edema, syncope, dysregulated body temperature, flank

pain, hypersensitivity reaction. Redistribution/accumulation of body fat (see [WARNINGS AND PRECAUTIONS, Fat Redistribution](#)).

Cardiovascular: arrhythmia, flushing, palpitations, tachycardia, thrombophlebitis, hypertension, congestive heart failure, chest pain

Central and Peripheral Nervous System: ataxia, confusion, convulsions, impaired coordination, migraine headaches, neuralgia, paresthesia, hypoesthesia, peripheral neuropathy, speech disorder, stupor, tremor, neuromuscular paresis, paranoid reaction

Gastrointestinal: dry mouth, pancreatitis, constipation, malabsorption

Liver and Biliary System: hepatic enzymes increased (including ALT, AST and GGT), hepatitis, jaundice, hepatomegaly (see [Post-Market Adverse Drug Reactions](#))

Metabolic and Nutritional: hypercholesterolemia, hypertriglyceridemia

Miscellaneous: thrombocytopenia, proteinuria, anemia, pancytopenia, increased sweating

Musculoskeletal: arthralgia, myalgia, myopathy, involuntary muscle contraction, muscle weakness, polyarthritis

Psychiatric: aggressive reactions, abnormal thinking, aggravated depression, agitation, delusions, amnesia, anxiety, apathy, delirium, depersonalization, emotional lability, euphoria, hallucination, manic reaction, psychosis, neurosis, paranoia, suicide, catatonia

Respiratory: asthma, apnea, dyspnea

Skin and Appendages: acne, alopecia, eczema, folliculitis, skin exfoliation, urticaria, erythema nodosum, erythema multiforme, Stevens-Johnson syndrome, verruca, nail disorders, skin disorders, photosensitivity reaction

Special Senses: abnormal vision, diplopia, glaucoma, iritis, parosmia, taste perversion, tinnitus

Urinary System: polyuria

### Abnormal Hematologic and Clinical Chemistry Findings

Table 5 summarizes clinically important laboratory abnormalities reported in Study 006 and ACTG 364.

**Table 5**  
**Selected Grade 3-4 Laboratory Abnormalities Reported in ≥2% of SUSTIVA-Treated Patients in Studies 006 and ACTG 364**

		Study 006 3TC-, NNRTI-, and Protease Inhibitor-Naive Patients			Study ACTG 364 NRTI-experienced, NNRTI- and Protease Inhibitor-Naive Patients		
Variable	Limit	SUSTIVA <sup>a</sup> + ZDV/3TC (n=412) 180 weeks <sup>b</sup>	SUSTIVA <sup>a</sup> + Indinavir (n=415) 102 weeks <sup>b</sup>	Indinavir + ZDV/3TC (n=401) 76 weeks <sup>b</sup>	SUSTIVA <sup>a</sup> + Nelfinavir + NRTIs (n=64) 71.1 weeks <sup>b</sup>	SUSTIVA <sup>a</sup> + NRTIs (n=65) 70.9 weeks <sup>b</sup>	Nelfinavir + NRTIs (n=66) 62.7 weeks <sup>b</sup>

Chemistry							
ALT	>5 x ULN	5%	8%	5%	2%	6%	3%
AST	>5 x ULN	5%	6%	5%	6%	8%	8%
GGT <sup>c</sup>	>5 x ULN	8%	7%	3%	5%	0	5%
Amylase	>2 x ULN	4%	4%	1%	0	6%	2%
Glucose	>250 mg/dL	3%	3%	3%	5%	2%	3%
Triglycerides <sup>d</sup>	≥751 mg/dL	9%	6%	6%	11%	8%	17%
Hematology							
Neutrophils	<750/mm <sup>3</sup>	10%	3%	5%	2%	3%	2%

a SUSTIVA provided as 600 mg once daily.

b Median duration of treatment.

c Isolated elevations of GGT in patients receiving SUSTIVA may reflect enzyme induction not associated with liver toxicity.

d Nonfasting.

ZDV = zidovudine, 3TC = lamivudine, ULN = Upper limit of normal, ALT = alanine aminotransferase,

AST = aspartate aminotransferase, GGT = gamma-glutamyltransferase.

**Liver Enzymes:** Liver function should be monitored in patients with a prior history of Hepatitis B and/or C (see [WARNINGS AND PRECAUTIONS; Monitoring and Laboratory Tests](#)).

In the long-term data set from Study 006, 137 patients treated with SUSTIVA-containing regimens (median duration of therapy, 68 weeks) and 84 treated with a control regimen (median duration, 56 weeks) were seropositive at screening for hepatitis B (surface antigen positive) and/or C (hepatitis C antibody positive). Among these co-infected patients, elevations in AST to greater than five times ULN developed in 13% of patients in the SUSTIVA arms and 7% of those in the control arm, and elevations in ALT to greater than five times ULN developed in 20% of patients in the SUSTIVA arms and 7% of patients in the control arm. Among co-infected patients, 3% of those treated with SUSTIVA-containing regimens and 2% in the control arm discontinued from the study because of liver or biliary system disorders.

**Lipids:** Increases in total cholesterol of 10-20% have been observed in some uninfected volunteers receiving SUSTIVA. In patients treated with SUSTIVA+ZDV+3TC, increases in non-fasting total cholesterol and HDL of approximately 20% and 25%, respectively, were observed. In patients treated with SUSTIVA+IDV, increases in non-fasting cholesterol and HDL of approximately 40% and 35%, respectively, were observed. Nonfasting total cholesterol levels ≥ 6.2 mmol/L and ≥ 7.8 mmol/L were reported in 34% and 9%, respectively, of patients treated with SUSTIVA + ZDV + 3TC; 54% and 20%, respectively, of patients treated with SUSTIVA + indinavir; and 28% and 4%, respectively, of patients treated with indinavir + ZDV + 3TC. The effects of SUSTIVA on triglycerides and LDL were not well-characterized since samples were taken from non-fasting patients. The clinical significance of these findings is unknown (see [WARNINGS AND PRECAUTIONS; Monitoring and Laboratory Tests](#)).

**Serum Amylase:** Asymptomatic elevations in serum amylase greater than 1.5 times the upper limit of normal were seen in 10% of patients treated with SUSTIVA and in 6% of patients treated with control regimens. The clinical significance of asymptomatic increases in serum amylase is unknown (see [WARNINGS AND PRECAUTIONS; Hepatic/Biliary/Pancreatic](#)).

## Post-Market Adverse Drug Reactions

Additional undesirable effects reported in postmarketing surveillance include neurosis, gynecomastia, rhabdomyolysis, increased CPK, blurred vision, photoallergic dermatitis, immune reconstitution inflammatory syndrome, cerebellar coordination and balance disturbances and vertigo.

Hepatic failure has been reported postmarketing, including some cases in patients with no pre-existing hepatic disease or other identifiable risk factors and also some cases characterized by a fulminant course, sometimes progressing to transplantation or death.

Additional cases of pancreatitis have been reported in postmarketing surveillance. Please see [WARNINGS AND PRECAUTIONS: Hepatic/Biliary/Pancreatic](#).

## DRUG INTERACTIONS

### Overview

Efavirenz has been shown *in vivo* to induce CYP3A4 and CYP2B6. Other compounds that are substrates of CYP3A4 or CYP2B6 may have decreased plasma concentrations when coadministered with SUSTIVA (efavirenz). *In vitro* studies have demonstrated that efavirenz inhibits 2C9, 2C19 and 3A4 isozymes in the range of observed efavirenz plasma concentrations. Coadministration of efavirenz with drugs primarily metabolized by these isozymes may result in altered plasma concentrations of the coadministered drug. Therefore, appropriate dose adjustments may be necessary for these drugs.

### Drug-Drug Interactions

Drugs which induce CYP3A4 activity (e.g., phenobarbital, rifampin, rifabutin) would be expected to increase the clearance of efavirenz resulting in lowered plasma concentrations. Drug interactions with SUSTIVA are summarized in Tables 1, 6 and 7. (See also [PHARMACOKINETICS; Drug-Drug Interactions](#) and [CONTRAINDICATIONS](#).)

There is limited information available on the potential for a pharmacodynamic interaction between SUSTIVA and drugs that prolong the QTc interval. QTc prolongation has been observed with the use of efavirenz (see [ACTION AND CLINICAL PHARMACOLOGY](#)). Consider alternatives to SUSTIVA when coadministered with a drug with a known risk of Torsade de Pointes.

The drugs listed in these tables are based on either drug interaction case reports or studies, or potential interactions due to the expected magnitude and seriousness of the interaction.

**Table 6**

Established Drug Interactions		
Concomitant Drug Class: Drug Name	Effect on Concentration	Clinical Comment
<i>Antiretroviral agents</i>		

Established Drug Interactions		
Concomitant Drug Class: Drug Name	Effect on Concentration	Clinical Comment
Protease inhibitor : Atazanavir	↓ atazanavir <sup>a</sup>	SUSTIVA decreases atazanavir exposure (see Table 9, ACTION AND CLINICAL PHARMACOLOGY, Drug-Drug Interactions).  <b>For treatment-naïve patients:</b> If atazanavir is combined with SUSTIVA, atazanavir 400 mg with ritonavir 100 mg should be administered once daily all as a single dose with food, and SUSTIVA should be administered on an empty stomach, preferably at bedtime.  <b>For treatment-experienced patients:</b> Do not coadminister atazanavir with SUSTIVA in treatment-experienced patients due to decreased atazanavir exposure.
Protease inhibitor : Fosamprenavir calcium	↓ amprenavir	For coadministration with fosamprenavir and ritonavir, the complete prescribing information for fosamprenavir calcium should be consulted.
Protease inhibitor : Indinavir	↓ indinavir <sup>a</sup>	The optimal dose of indinavir, when given in combination with SUSTIVA, is not known. Increasing the indinavir dose to 1000 mg every 8 hours does not compensate for the increased indinavir metabolism due to SUSTIVA. When indinavir at an increased dose (1000 mg every 8 hours) was given with SUSTIVA (600 mg once daily), the indinavir AUC and C <sub>min</sub> were decreased on average by 33%-46% and 39-57%, respectively, compared to when indinavir (800 mg every 8 hours) was given alone.
Protease inhibitor : Lopinavir/ritonavir	↓ lopinavir <sup>a</sup>	For lopinavir/ritonavir capsules or oral solution, a dose increase to 533/133 mg (4 capsules or 6.5 mL) twice daily taken with food is recommended when used in combination with SUSTIVA in patients for whom reduced susceptibility to lopinavir is clinically suspected (by treatment history or laboratory evidence). <b>Lopinavir/ritonavir tablets should not be administered once-daily in combination with SUSTIVA.</b>  In antiretroviral-naïve patients, lopinavir/ritonavir tablets can be used twice daily in combination with SUSTIVA with no dose adjustment. A dose increase of lopinavir/ritonavir tablets to 600/150 mg (3 tablets) twice daily may be considered when used in combination with SUSTIVA in treatment-experienced patients where decreased susceptibility to lopinavir is clinically suspected (by treatment history or laboratory evidence).
CCR5 co-receptor antagonist : Maraviroc	↓ maraviroc <sup>a</sup>	Refer to the full prescribing information for maraviroc for guidance on coadministration with efavirenz.
Integrase strand transfer inhibitor: Raltegravir	↓ raltegravir <sup>a</sup>	Efavirenz did not have a clinically meaningful effect on the pharmacokinetics of raltegravir.

Established Drug Interactions		
Concomitant Drug Class: Drug Name	Effect on Concentration	Clinical Comment
Protease inhibitor : Ritonavir	↑ritonavir <sup>a</sup> ↑efavirenz <sup>a</sup>	When ritonavir 500 mg q12h was coadministered with SUSTIVA 600 mg once daily, the combination was associated with a higher frequency of adverse clinical experiences (e.g., dizziness, nausea, paresthesia) and laboratory abnormalities (elevated liver enzymes). Monitoring of liver enzymes is recommended when SUSTIVA is used in combination with ritonavir.
Protease inhibitor : Saquinavir	↓ saquinavir <sup>a</sup>	Should not be used as sole protease inhibitor in combination with SUSTIVA.
<i>Hepatitis C antiviral agents</i>		
NS5B polymerase inhibitor /NS5A inhibitor: Sofosbuvir/ velpatasvir	↔sofosbuvir ↓velpatasvir ↔efavirenz	Coadministration of sofosbuvir/velpatasvir with efavirenz is not recommended. Concomitant administration of sofosbuvir/velpatasvir with efavirenz resulted in a reduction (approximately 50%) in the systemic exposure of velpatasvir. The mechanism of the effect on velpatasvir is induction of CYP3A and CYP2B6 by efavirenz. Refer to the prescribing information for sofosbuvir/velpatasvir for more information.
Velpatasvir/ sofosbuvir/ voxilaprevir	↓velpatasvir ↓voxilaprevir	Concomitant administration of velpatasvir/sofosbuvir/ voxilaprevir with efavirenz is not recommended, as it may decrease concentrations of velpatasvir and voxilaprevir. Refer to the prescribing information for velpatasvir/sofosbuvir/ voxilaprevir for more information.
Elbasvir/ grazoprevir	↓elbasvir ↓grazoprevir ↔efavirenz	Concomitant administration of efavirenz with elbasvir/grazoprevir is contraindicated because it may lead to loss of virologic response to elbasvir/grazoprevir. This loss is due to significant decreases in elbasvir and grazoprevir plasma concentrations caused by CYP3A4 induction. Refer to the prescribing information for elbasvir/grazoprevir for more information.
Protease inhibitor : Glecaprevir/ pibrentasvir	↓glecaprevir ↓pibrentasvir ↔efavirenz	Concomitant administration of glecaprevir/pibrentasvir with SUSTIVA may significantly decrease plasma concentrations of glecaprevir and pibrentasvir, leading to reduced therapeutic effect. Coadministration of glecaprevir/pibrentasvir with SUSTIVA is not recommended. Refer to the prescribing information for glecaprevir/ pibrentasvir for more information.
<i>Other agents</i>		
Anticonvulsants: Carbamazepine	↓ carbamazepine <sup>a</sup> ↓ efavirenz <sup>a</sup>	Plasma concentrations of carbamazepine and SUSTIVA decreased. Periodic monitoring of carbamazepine plasma levels should be conducted. There are insufficient data to make a dose recommendation. Alternative anticonvulsant treatment should be considered.

Established Drug Interactions		
Concomitant Drug Class: Drug Name	Effect on Concentration	Clinical Comment
Antidepressants: Bupropion	↓ bupropion <sup>a</sup>	The effect of efavirenz on bupropion exposure is thought to be due to the induction of bupropion metabolism. Bupropion dose adjustments should be guided by clinical response, but the maximum recommended dose of bupropion should not be exceeded. No dose adjustment is necessary for efavirenz.
Sertraline	↓ sertraline <sup>a</sup>	Since SUSTIVA reduces sertraline levels, it may be necessary to retitrate the sertraline dose in order to achieve the desired clinical effect. In a drug interaction study in healthy subjects, an increased incidence of impaired concentration was seen in subjects receiving sertraline concomitantly with SUSTIVA.
Antifungal: Itraconazole	↓ itraconazole <sup>a</sup> ↓ hydroxyitraconazole <sup>a</sup>	Since no dose recommendation for itraconazole can be made, alternative antifungal treatment should be considered.
Antifungal: Posaconazole	↓ posaconazole <sup>a</sup>	Avoid concomitant use of posaconazole and efavirenz unless the benefit to the patient outweighs the risk.
Antifungal: Voriconazole	↓ voriconazole <sup>a</sup> ↑ efavirenz <sup>a</sup>	Standard doses of voriconazole and SUSTIVA should not be used concurrently. When voriconazole is coadministered with SUSTIVA, the voriconazole maintenance dose should be increased to 400 mg every 12 hours and SUSTIVA dose should be decreased to 300 mg once daily.
Anthelmintic: Praziquantel	↓ praziquantel	Coadministration with efavirenz is not recommended due to significant decrease in plasma concentrations of praziquantel, with risk of treatment failure due to increased hepatic metabolism by efavirenz. In case the combination is needed, an increased dose of praziquantel could be considered.
Anti-infective: Clarithromycin	↓ clarithromycin <sup>a</sup> ↑ 14-OH metabolite <sup>a</sup>	Consider alternatives to macrolide antibiotics because of the risk of QT interval prolongation. Plasma concentrations decreased by SUSTIVA; clinical significance unknown. In uninfected volunteers, 46% developed rash while receiving SUSTIVA and clarithromycin. No dose adjustment of SUSTIVA is recommended when given with clarithromycin. Alternatives to clarithromycin, such as azithromycin, should be considered (see section on “Other Drugs”, below Table 7). Not all macrolide antibiotics have been studied in combination with SUSTIVA.
Antimalarial: Artemether/ lumefantrine <sup>b</sup> :	↓ artemether <sup>a</sup> ↓ dihydroartemisinin <sup>a</sup> ↓ lumefantrine <sup>a</sup>	Consider alternatives to artemether/lumefantrine because of the risk of QT interval prolongation. Coadministration of efavirenz with artemether /lumefantrine resulted in a decrease in exposures to artemether, dihydroartemisinin (active metabolite of artemether), and lumefantrine. Exposure to efavirenz was not significantly affected. Since decreased concentrations of artemether, dihydroartemisinin, or lumefantrine may result in a decrease of antimalarial efficacy, caution is recommended when SUSTIVA and artemether/lumefantrine tablets are coadministered.

Established Drug Interactions		
Concomitant Drug Class: Drug Name	Effect on Concentration	Clinical Comment
Antimalarial: Atovaquone Proguanil	↓ atovaquone ↓ proguanil	Coadministration of efavirenz with atovaquone/ proguanil resulted in a decrease in exposures to atovaquone and proguanil. Since decreased concentrations of atovaquone and proguanil may result in a decrease of antimalarial efficacy, concomitant administration should be avoided whenever possible.
Antimycobacterial: Rifabutin	↓ rifabutin <sup>a</sup>	Consider an increase of the daily dose of rifabutin by 50%. Consider doubling the rifabutin dose in regimens where rifabutin is given 2 or 3 times a week.
Antimycobacterial: Rifampin	↓ efavirenz <sup>a</sup>	Rifampin has the potential to decrease serum concentration of SUSTIVA. When efavirenz is coadministered with rifampin to adult patients weighing 50 kg or more, a dose increase of efavirenz to 800 mg once daily is recommended.
Calcium channel blocker: Diltiazem	↓ diltiazem <sup>a</sup> ↓ desacetyl diltiazem <sup>a</sup> ↓ N-monodesmethyl diltiazem <sup>a</sup>	Diltiazem levels are markedly decreased when coadministered with SUSTIVA. SUSTIVA levels increased to a lesser extent (see Tables 9 and 10). Patients should be closely monitored for possible decreased diltiazem effects and increased adverse events and laboratory abnormalities associated with SUSTIVA. Refer to the prescribing information for diltiazem for guidance on dose adjustment).
HMG-CoA Reductase Inhibitors: Atorvastatin Pravastatin Simvastatin	↓ atorvastatin <sup>a</sup> ↓ pravastatin <sup>a</sup> ↓ simvastatin <sup>a</sup>	Plasma concentrations of atorvastatin and pravastatin decreased. Consult the complete prescribing information for the HMG-CoA reductase inhibitor for guidance on individualizing the dose. A marked decrease in simvastatin plasma concentrations was seen when co-administered with SUSTIVA (see Table 9). Alternative statins should be considered.
Hormonal contraceptive:  Oral: Ethinyl estradiol/ Norgestimate  Implant: Etonogestrel	↓ active metabolites of norgestimate <sup>a</sup>  ↓ etonogestrel	A reliable method of barrier contraception must be used in addition to hormonal contraceptives.  Efavirenz had no effect on ethinyl estradiol concentrations, but progestin levels (norelgestromin and levonorgestrel) were markedly decreased. The clinical significance of these effects is not known. No effect of ethinyl estradiol/norgestimate on efavirenz plasma concentrations was observed.  Decreased exposure of etonogestrel may be expected (CYP3A4 induction), and there have been post-marketing reports of contraceptive failure with etonogestrel in efavirenz-exposed patients.
Narcotic analgesic: Methadone	↓ methadone <sup>a</sup>	Coadministration in HIV-infected individuals with a history of injection drug use resulted in decreased plasma levels of methadone and signs of opiate withdrawal. Patients should be monitored for signs of withdrawal and their methadone dose increased as required to alleviate withdrawal symptoms.

<sup>a</sup> For magnitude of interactions, see Tables 9 and 10.

<sup>b</sup> Not marketed in Canada

**Table 7**

<b>Other Potentially Clinically Significant Drug Interactions With SUSTIVA<sup>a</sup></b>	
Anticoagulants: Warfarin Acenocoumarol	Plasma concentrations and effects potentially increased or decreased by SUSTIVA. It is recommended that INR be monitored.
Anticonvulsants: Phenytoin Phenobarbital	Potential for reduction in anticonvulsant and/or efavirenz plasma levels; periodic monitoring of anticonvulsant plasma levels should be conducted. (See WARNINGS AND PRECAUTIONS.)
Antifungals: Ketoconazole	See CONTRAINDICATIONS for other antifungals. Drug interaction studies with SUSTIVA and ketoconazole have not been conducted. SUSTIVA has the potential to decrease plasma concentrations of ketoconazole.
Anti-HIV protease inhibitors: Saquinavir/ritonavir combination	No pharmacokinetic data are available. (See Table 6.)
Calcium channel blockers: felodipine, nifedipine, verapamil	No data are available on the potential interactions of efavirenz with calcium channel blockers that are substrates of the CYP3A4 enzyme, other than diltiazem (see Table 6). The potential exists for reduction in plasma concentrations of the calcium channel blocker. Dose adjustments should be guided by clinical response (refer to the prescribing information for the calcium channel blocker).
Immunosuppressants: cyclosporine, tacrolimus, sirolimus	When an immunosuppressant metabolized by CYP3A4 is administered with efavirenz, decreased exposure of the immunosuppressant may be expected due to CYP3A4 induction. Dose adjustments of the immunosuppressant may be required. Close monitoring of immunosuppressant concentrations for at least 2 weeks (until stable concentrations are reached) is recommended when starting or stopping treatment with SUSTIVA.
Non-nucleoside reverse transcriptase inhibitors	No studies have been performed with other NNRTIs.

<sup>a</sup> This table is not all inclusive.

**Other Drugs:** Based on the results of drug interaction studies, no dosage adjustment of either SUSTIVA or the following coadministered drugs is recommended: aluminum/magnesium hydroxide antacids, azithromycin, cetirizine, famotidine, fluconazole, lamivudine, nelfinavir, paroxetine, zidovudine and tenofovir disoproxil fumarate. (See [ACTION AND CLINICAL PHARMACOLOGY](#); [Pharmacokinetics](#), [Tables 9](#) and [10](#).)

No dosage adjustment for lorazepam is recommended when coadministered with SUSTIVA.

Specific drug interaction studies have not been performed with SUSTIVA and NRTIs other than lamivudine and zidovudine. Clinically significant interactions would not be expected since the NRTIs are metabolized via a different route than efavirenz and would be unlikely to compete for the same metabolic enzymes and elimination pathways.

### **Drug-Food Interactions**

Food increases the concentrations of efavirenz which may lead to an increase in frequency of adverse events. Therefore, it is recommended that SUSTIVA be taken on an empty stomach (See [DOSAGE AND ADMINISTRATION](#); [Recommended Dose and Dosage Adjustment](#) and

[ACTION AND CLINICAL PHARMACOLOGY; Pharmacokinetics, Effect of Food on Oral Absorption](#)).

### **Drug-Herb Interactions**

**St. John's Wort:** Concomitant use of St. John's wort (*Hypericum perforatum*) or St. John's wort-containing products with efavirenz is contraindicated. Coadministration of non-nucleoside reverse transcriptase inhibitors (NNRTIs), including SUSTIVA, with St. John's wort is expected to substantially decrease NNRTI concentrations. Decreased concentrations may result in suboptimal levels of efavirenz and lead to loss of virologic response and possible resistance to efavirenz or to the class of NNRTIs.

### **Drug-Laboratory Test Interactions**

**Cannabinoid Test Interaction:** Efavirenz does not bind to cannabinoid receptors. False-positive urine cannabinoid test results have been reported with some screening assays in uninfected and HIV-infected subjects receiving efavirenz. Confirmation of positive screening tests for cannabinoids by a more specific method such as gas chromatography/mass spectrometry is recommended.

## **DOSAGE AND ADMINISTRATION**

### **Dosing Considerations**

SUSTIVA must be given in combination with other antiretroviral medications.

### **Recommended Dose and Dosage Adjustment**

#### **Adults**

The recommended dosage of SUSTIVA (efavirenz) capsules in combination with other antiretroviral agents is 600 mg orally, once daily. It is recommended that SUSTIVA be taken on an empty stomach, preferably at bedtime. The increased efavirenz concentrations observed following administration of SUSTIVA with food may lead to an increase in frequency of adverse events (see [ACTIONS AND CLINICAL PHARMACOLOGY; Effect of Food on Oral Absorption](#)). Dosing at bedtime may improve the tolerability of nervous system symptoms (see [WARNINGS AND PRECAUTIONS; Neurologic](#) and [ADVERSE REACTIONS](#)).

#### **Pediatric Patients and Adolescents**

It is recommended that SUSTIVA be taken on an empty stomach, preferably at bedtime. The recommended dosage of SUSTIVA in combination with other antiretroviral agents for patients 3 to 17 years of age is described in Table 8. There are insufficient data to recommend a dose in pediatric patients below 3 years of age or who weigh less than 13 kg. SUSTIVA capsules must only be administered to children who are able to reliably swallow capsules. The recommended dosage of SUSTIVA for pediatric patients weighing greater than 40 kg is 600 mg, once daily.

**Table 8**  
**Pediatric Dose to be Administered Once Daily**

Body Weight		SUSTIVA Dose (mg)
kg	lbs	
13 to < 15	29 to < 33	200
15 to < 20	33 to < 44	250
20 to < 25	44 to < 55	300
25 to < 32.5	55 to < 71.5	350
32.5 to < 40	71.5 to < 88	400
≥ 40	≥ 88	600

Information is based on one study, ACTG 382. Patients were administered SUSTIVA in combination with nelfinavir and NRTIs.

Further dose adjustments may be required if other products are used concomitantly. (See [Pharmacokinetics: Drug-Drug Interactions](#) and [DRUG INTERACTIONS](#).)

#### [Renal Impairment](#)

See [WARNINGS AND PRECAUTIONS](#); [Special Populations](#), [Renal Impairment](#).

#### [Hepatic Impairment](#)

SUSTIVA use is not recommended in patients with moderate or severe hepatic impairment (Child-Pugh Class B or C) because of insufficient data to determine appropriate dosing. Caution should be exercised in administering SUSTIVA to patients with mild hepatic impairment because of the extensive cytochrome P450-mediated metabolism of efavirenz and limited clinical experience. See [WARNINGS AND PRECAUTIONS](#); [Special Populations](#), [Hepatic Impairment, Monitoring and Laboratory Tests](#), [Liver enzymes](#) and [ACTION AND CLINICAL PHARMACOLOGY](#); [Pharmacokinetics](#); [Special Populations](#).

#### **Missed Dose**

If a dose of SUSTIVA is missed, patients should take the dose as soon as possible and then return to their normal schedule. However, if a dose is skipped, the patient should not double the next dose.

#### **OVERDOSAGE**

For management of a suspected drug overdose, please contact your regional Poison Control Center.

Treatment of overdose with SUSTIVA (efavirenz) should consist of general supportive measures, including monitoring of vital signs and observation of the patient's clinical status. Administration of activated charcoal should be used to aid removal of unabsorbed drug, as recommended in American College of Emergency Physicians guidelines. There is no specific

antidote for overdose with SUSTIVA. Since efavirenz is highly protein bound, dialysis is unlikely to significantly remove the drug from blood.

Some patients accidentally taking 600 mg twice daily have reported increased nervous system symptoms. One patient experienced involuntary muscle contractions and a second patient experienced vomiting after taking twice the recommended dose.

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

Efavirenz is a selective non-nucleoside reverse transcriptase (RT) inhibitor of human immunodeficiency virus type 1 (HIV-1). Efavirenz is predominantly a non-competitive inhibitor of HIV-1 RT. HIV-2 RT and human cellular DNA polymerases  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$  are not inhibited by concentrations of efavirenz well in excess of those achieved clinically (see [VIROLOGY](#)).

### **Pharmacodynamics**

Cardiac Electrophysiology: The effect of SUSTIVA on the QTc interval was evaluated in an open-label, positive and placebo controlled, fixed single sequence 3-period, 3-treatment crossover QT study in 58 healthy subjects enriched for CYP2B6 polymorphisms. The mean  $C_{max}$  of efavirenz in subjects with CYP2B6 \*6/\*6 genotype following the administration of 600 mg daily dose for 14 days was 2.25-fold the mean  $C_{max}$  observed in subjects with CYP2B6 \*1/\*1 genotype. A positive relationship between efavirenz concentration and QTc prolongation was observed. Based on the concentration-QTc relationship, the mean QTc prolongation and its upper bound 90% confidence interval are 8.7 ms and 11.3 ms in subjects with CYP2B6\*6/\*6 genotype following the administration of 600 mg daily dose for 14 days (see [WARNING AND PRECAUTION](#)).

### **Pharmacokinetics**

#### Absorption

Peak efavirenz plasma concentrations were attained within 5 hours following single oral doses of 100 mg to 1600 mg administered to uninfected volunteers. Dose-related increases in  $C_{max}$  and AUC were seen for doses up to 1600 mg.

In HIV-infected patients at steady-state, mean  $C_{max}$ , mean  $C_{min}$ , and mean AUC were dose proportional.

#### Effect of Food on Oral Absorption

Administration of a single 600 mg dose of efavirenz capsules with a high fat/high caloric meal (894 kcal, 54 g fat, 54% calories from fat) or a reduced fat/normal caloric meal (440 kcal, 2 g fat, 4% calories from fat) was associated with a mean increase of 22% and 17% in efavirenz AUC and a mean increase of 39% and 51% in efavirenz  $C_{max}$ , respectively, relative to the exposures

achieved when given under fasted conditions (see [DOSAGE AND ADMINISTRATION; Recommended Dose and Dosage Adjustment](#)).

### Distribution

Efavirenz is highly bound (approximately 99.5-99.75%) to human plasma proteins, predominantly albumin. In HIV-1 infected patients (N=9), efavirenz cerebrospinal fluid concentrations ranged from 0.26 to 1.19% (mean 0.69%) of the corresponding plasma concentration; approximately 3-fold higher than the non-protein-bound (free) fraction of efavirenz in plasma.

### Metabolism

*In vivo* and *in vitro* studies demonstrated that efavirenz is principally metabolized by the cytochrome P450 system to hydroxylated metabolites with subsequent glucuronidation of these hydroxylated metabolites. These metabolites are essentially inactive against HIV-1.

Efavirenz has been shown to induce P450 enzymes, resulting in the induction of its own metabolism.

### Elimination

Efavirenz has a long terminal half-life of 52-76 hours after single doses and 40-55 hours after multiple doses. Approximately 14-34% of the radiolabel was recovered in the urine and 16-61% was recovered in the feces. Nearly all of the urinary excretion of the radiolabelled drug was in the form of metabolites. Efavirenz accounted for the majority of the total radioactivity measured in feces (see [PHARMACOLOGY; Pharmacokinetics](#) for more detail.)

### Special Populations

#### Hepatic Impairment

In a multiple-dose SUSTIVA PK study (600 mg daily), the mean C<sub>max</sub> and mean AUC of efavirenz in patients with mild hepatic impairment (Child-Pugh Class A, n=6) were 20.3±15.5 µM (mean±SD) and 351±336.9 µM•h (mean±SD), respectively, compared with those in controls (n=6; C<sub>max</sub>=28.4±27.35 µM, AUC=506±581 µM•h). There were insufficient data to determine whether moderate or severe hepatic impairment (Child-Pugh Class B or C) affects efavirenz pharmacokinetics (see [WARNINGS AND PRECAUTIONS](#) and [DOSAGE AND ADMINISTRATION](#)).

### Drug-Drug Interactions

Tables 9 and 10 show drug-drug interactions of SUSTIVA with various co-administered drugs and their pharmacokinetic profiles (see [DRUG INTERACTIONS](#)).

**Table 9**  
**Effect of Efavirenz on Coadministered Drug Plasma C<sub>max</sub>, AUC and C<sub>min</sub>**

Coadministered Drug Class: Drug Name	Dose	Efavirenz Dose	Coadministered Drug (Mean % change)		
			Cmax	AUC	Cmin
<i>Antiretroviral agents</i>					
Protease inhibitor : Atazanavir	400 mg daily x 20 days	600 mg d 7-20	↓ 59%	↓ 74%	↓ 93%
Atazanavir/ritonavir	400 mg daily d 1-6, then 300 mg daily d 7-20 with ritonavir 100 mg daily and a light meal	600 mg daily 2 h after atazanavir and ritonavir d 7-20	↑ 14% <sup>a</sup>	↑ 39% <sup>a</sup>	↑ 48% <sup>a</sup>
	300 mg qd/ritonavir 100 mg qd d 1-10 (pm), then 400 mg qd/ritonavir 100 mg qd d 11-24 (pm) (simultaneous with efavirenz)	600 mg d 11-24 (pm)	↑ 17%	↔	↓ 42%
Protease inhibitor : Indinavir	1000 mg q8h x 10 days after morning dose after afternoon dose after evening dose	600 mg x 10 days	↔ <sup>b</sup> ↔ <sup>b</sup> ↓ 29% <sup>b</sup>	↓ 33% <sup>b</sup> ↓ 37% <sup>b</sup> ↓ 46% <sup>b</sup>	↓ 39% <sup>b</sup> ↓ 52% <sup>b</sup> ↓ 57% <sup>b</sup>
Protease inhibitor : Indinavir / ritonavir	Indinavir 800 mg + ritonavir 100 mg q12h d 1-29	600 mg d 15-29	↓ 17% <sup>e</sup>	↓ 25% <sup>e</sup>	↓ 50% <sup>e</sup>
Protease inhibitor : Lopinavir/ritonavir	400/100 mg capsule q12h x 9 days	600 mg x 9 days	↔ <sup>c</sup>	↓ 19% <sup>c</sup>	↓ 39% <sup>c</sup>
	600/150 mg tablet q12h x 10 days with efavirenz compared to 400/100 mg q12h alone	600 mg x 9 days	↑ 36%	↑ 36%	↑ 32%
CCR5 co-receptor antagonist: Maraviroc	100 mg bid	600 mg	↓ 51%	↓ 45%	↓ 45%
Integrase strand transfer inhibitor: Raltegravir	400 mg single dose	600 mg	↓ 36%	↓ 36%	↓ 21%
Protease inhibitor : Nelfinavir Metabolite AG-1402	750 mg q8h x 7 days	600 mg x 7 days	↑ 21% ↓ 40%	↑ 20% ↓ 37%	↔ ↓ 43%

Coadministered Drug Class: Drug Name	Dose	Efavirenz Dose	Coadministered Drug (Mean % change)		
			Cmax	AUC	Cmin
Protease inhibitor : Ritonavir	500 mg q12h x 8 days after AM dose after PM dose	600 mg x 10 days	↑ 24% ↔	↑ 18% ↔	↑ 42% ↑ 24%
Protease inhibitor : Saquinavir (SGC) <sup>g</sup>	1200 mg q8h x 10 days	600 mg x 10 days	↓ 50%	↓ 62%	↓ 56%
Nucleoside reverse transcriptase inhibitor : Lamivudine	150 mg q12h x 14 days	600 mg x 14 days	↔	↔	↑ 265%
Nucleoside reverse transcriptase inhibitor : Zidovudine	300 mg q12h x 14 days	600 mg x 14 days	↔	↔	↑ 225%
Nucleotide reverse transcriptase inhibitor: Tenofovir disoproxil fumarate	300 mg daily	600 mg x 14 days	↔	↔	↔
<i>Hepatitis C antiviral agents</i>					
NS5B polymerase inhibitor /NS5A inhibitor: Sofosbuvir/ velpatasvir <sup>i</sup>	400 mg once daily/ 100 mg once daily	600 mg once daily	↑ 38%/ ↓ 47%	↔/ ↓ 53%	NA/ ↓ 57%
<i>Other agents</i>					
Anticonvulsant: Carbamazepine	200 mg daily x 3 days, 200 mg bid x 3 days, then 400 mg daily x 29 days	600 mg x 14 days	↓ 20%	↓ 27%	↓ 35%
Epoxide metabolite			↔	↔	↓ 13%
Antidepressant: Bupropion	150 mg single dose (sustained-release)	600 mg x 14 days	↓ 34%	↓ 55%	NA
Hydroxybupropion			↑ 50%	↔	NA
Antidepressant: Paroxetine	20 mg daily x 14 days	600 mg x 14 days	↔	↔	↔
Antidepressant: Sertraline N desmethylsertraline	50 mg daily x 14 days	600 mg x 14 days	↓ 29% ↓ 17%	↓ 39% ↓ 20%	↓ 46% ↓ 20%
Antifungal: Fluconazole	200 mg x 7 days	400 mg x 7 days	↔	↔	↔

Coadministered Drug Class: Drug Name	Dose	Efavirenz Dose	Coadministered Drug (Mean % change)		
			Cmax	AUC	Cmin
Antifungal: Itraconazole Hydroxyitraconazole	200 mg q12h x 28 days	600 mg x 14 days	↓ 37% ↓ 35%	↓ 39% ↓ 37%	↓ 44% ↓ 43%
Antifungal: Posaconazole	400 mg (oral suspension) bid x 10 and 20 days	400 mg x 10 and 20 days	↓ 45%	↓ 50%	NA
Antifungal: Voriconazole	400 mg q12h d -1 200 mg q12h d 2-9  300 mg po q12h d 2-7 400 mg po q12h d 2-7	400 mg x 9 days  300 mg x 7 days 300 mg x 7 days	↓ 61%  ↓ 36% <sup>f</sup> ↑ 23% <sup>f</sup>	↓ 77%  ↓ 55% <sup>f</sup> ↔ <sup>f</sup>	NA  NA NA
Anti-infective: Azithromycin	600 mg single dose	400 mg x 7 days	↑ 22%	↔	NA
Anti-infective: Clarithromycin 14-OH metabolite	500 mg q12h x 7 days	400 mg x 7 days	↓ 26% ↑ 49%	↓ 39% ↑ 34%	↓ 53% ↑ 26%
Antimalarial: Artemether <sup>h</sup>  dihydroartemisinin (active metabolite of artemether) <sup>h</sup>  lumefantrine <sup>h</sup>	80/480 mg bid x 3 days before and during efavirenz coadministration	600 mg x 26 days	↓ 21%  ↓ 38%  ↔	↓ 51%  ↓ 46%  ↓ 21%	NA  NA  NA
Antimycobacterial: Rifabutin 25-0- desacetyl rifabutin	300 mg daily x 14 days	600 mg x 14 days	↓ 32% ↓ 49% <sup>d</sup>	↓ 38% ↓ 74% <sup>d</sup>	↓ 45% NA
Anxiolytic: Lorazepam	2 mg single dose	600 mg x 10 days	↑ 16%	↔	NA
Calcium channel blocker: Diltiazem Desacetyl diltiazem N-monodesmethyl	240 mg x 21 days	600 mg x 14 days	↓ 60% ↓ 64% ↓ 28%	↓ 69% ↓ 75% ↓ 37%	↓ 63% ↓ 62% ↓ 37%
H <sub>1</sub> receptor antagonist: Cetirizine	10 mg single dose	600 mg x 10 days	↓ 24%	↔	NA

Coadministered Drug Class: Drug Name	Dose	Efavirenz Dose	Coadministered Drug (Mean % change)		
			C <sub>max</sub>	AUC	C <sub>min</sub>
HMG-CoA reductase inhibitor: Atorvastatin Total active (including metabolites)	10 mg daily x 4 days	600 mg x 15 days	↓ 14% ↓ 15%	↓ 43% ↓ 32%	↓ 69% ↓ 48%
HMG-CoA reductase inhibitor: Pravastatin	40 mg daily x 4 days	600 mg x 15 days	↓ 32%	↓ 44%	↓ 19%
HMG-CoA reductase inhibitor: Simvastatin Total active (including metabolites)	40 mg daily x 4 days	600 mg x 15 days	↓ 72% ↓ 68%	↓ 68% ↓ 60%	↓ 45% NA
Narcotic analgesic: Methadone	Stable maintenance 35-100 mg daily	600 mg x 14-21 days	↓ 45%	↓ 52%	NA
Oral contraceptive: Ethinyl estradiol/ Norgestimate  Ethinyl estradiol Norelgestromin Levonorgestrel	0.035 mg/ 0.25 mg x 14 days	600 mg x 14 days	↔ ↓ 46% ↓ 80%	↔ ↓ 64% ↓ 83%	↔ ↓ 82% ↓ 86%

↑ Indicates increase ↓ Indicates decrease ↔ Indicates no change or a mean increase or decrease of <10%

<sup>a</sup> Compared with atazanavir 400 mg daily alone.

<sup>b</sup> Comparator dose of indinavir was 800 mg q8h x 10 days.

<sup>c</sup> Values are for lopinavir. C<sub>min</sub> of lopinavir was significantly decreased by 39%. C<sub>max</sub> and AUC of lopinavir were decreased by 3% and 19% respectively (not significant). The pharmacokinetics of ritonavir 100 mg q12h are unaffected by concurrent efavirenz.

<sup>d</sup> Based on arithmetic mean values.

<sup>e</sup> Compared to indinavir 800 twice daily given with ritonavir 100 mg twice daily without efavirenz. The geometric C<sub>min</sub> for indinavir (0.33 mg/L) when given with ritonavir and efavirenz was higher than the mean historical C<sub>min</sub> (0.15 mg/L) when indinavir was given alone at 800 mg every 8 hours. When efavirenz 600 mg once daily was given with indinavir/ritonavir 800/100 mg twice daily in HIV-1-infected patients (n=6), the pharmacokinetics of indinavir and efavirenz were generally comparable to these data from uninfected volunteers.

<sup>f</sup> Relative to steady-state administration of voriconazole (400 mg for 1 day, then 200 mg po q12h for 2 days).

<sup>g</sup> SGC Soft Gelatin Capsule

<sup>h</sup> Not marketed in Canada

<sup>i</sup> Administered as ATRIPLA (efavirenz, emtricitabine and tenofovir DF fixed-dose combination)

**Table 10**  
**Effect of Coadministered Drug on Efavirenz Plasma C<sub>max</sub>, AUC and C<sub>min</sub>**

Coadministered Drug Class: Drug Name	Dose	Efavirenz Dose	Efavirenz (Mean % change)		
			C <sub>max</sub>	AUC	C <sub>min</sub>
<i>Antiretroviral agents</i>					
Protease inhibitor : Atazanavir	400 mg daily x 20 days	600 mg d 7-20	↔	↔	NA
Protease inhibitor : Indinavir	800 mg q8h x 14 days	200 mg x 14 days	↔	↔	↔
Protease inhibitor : Lopinavir/ritonavir	400/100 mg q12h x 9 days	600 mg x 9 days	↔	↓ 16%	↓ 16%
Protease inhibitor : Nelfinavir	750 mg q8h x 7 days	600 mg x 7 days	↓ 12%	↓ 12%	↓ 21%
Protease inhibitor : Ritonavir	500 mg q12h x 8 days	600 mg x 10 days	↑ 14%	↑ 21%	↑ 25%
Protease inhibitor : Saquinavir (SGC) <sup>b</sup>	1200 mg q8h x 10 days	600 mg x 10 days	↓ 13%	↓ 12%	↓ 14%
Nucleotide reverse transcriptase inhibitor: Tenofovir disoproxil fumarate	300 mg daily	600 mg x 14 days	↔	↔	↔
<i>Hepatitis C antiviral agents</i>					
NS5B polymerase inhibitor /NS5A inhibitor: Sofosbuvir /velpatasvir <sup>d</sup>	400 mg/100 mg daily	600 mg daily	↔	↔	↔
<i>Other agents</i>					
Antacid: Aluminum hydroxide 400 mg Magnesium hydroxide 400 mg, + simethicone 30 mg	30 mL single dose	400 mg single dose	↔	↔	NA
Anticonvulsant: Carbamazepine	200 mg daily x 3 days, 200 mg bid x 3 days, then 400 mg daily x 15 days	600 mg x 35 days	↓ 21%	↓ 36%	↓ 47%
Antidepressant: Paroxetine	20 mg daily x 14 days	600 mg x 14 days	↔	↔	↔
Antidepressant: Sertraline	50 mg daily x 14 days	600 mg x 14 days	↑ 11%	↔	↔
Antifungal: Fluconazole	200 mg x 7 days	400 mg x 7 days	↔	↑ 16%	↑ 22%
Antifungal: Itraconazole	200 mg q12h x 14 days	600 mg x 28 days	↔	↔	↔

Coadministered Drug Class: Drug Name	Dose	Efavirenz Dose	Efavirenz (Mean % change)		
			C <sub>max</sub>	AUC	C <sub>min</sub>
Antifungal: Voriconazole	400 mg q12h d-1	400 mg x 9 days	↑ 38%	↑ 44%	NA
	200 mg q12h d 2-9				
	300 mg po q12h d 2-7	300 mg x 7 days	↓ 14% <sup>a</sup>	↔ <sup>a</sup>	NA
	400 mg po q12h d 2-7	300 mg x 7 days	↔ <sup>a</sup>	↑ 17% <sup>a</sup>	NA
Anti-infective: Azithromycin	600 mg single dose	400 mg x 7 days	↔	↔	↔
Anti-infective: Clarithromycin	500 mg q12h x 7 days	400 mg x 7 days	↑ 11%	↔	↔
Antimycobacterial: Rifabutin	300 mg daily x 14 days	600 mg x 14 days	↔	↔	↓ 12%
Antimycobacterial: Rifampin	600 mg x 7 days	600 mg x 7 days	↓ 20%	↓ 26%	↓ 32%
Calcium channel blocker: Diltiazem	240 mg x 14 days	600 mg x 28 days	↑ 16%	↑ 11%	↑ 13%
H <sub>1</sub> receptor antagonist: Cetirizine	10 mg single dose	600 mg x 10 days	↔	↔	↔
H <sub>2</sub> receptor antagonist: Famotidine	40 mg single dose	400 mg single dose	↔	↔	NA
HMG-CoA reductase inhibitor: Atorvastatin	10 mg daily x 4 days	600 mg x 15 days	↔	↔	↔
HMG-CoA reductase inhibitor: Pravastatin	40 mg daily x 4 days	600 mg x 15 days	↔	↔	↔
HMG-CoA reductase inhibitor: Simvastatin	40 mg daily x 4 days	600 mg x 15 days	↓ 12%	↔	↓ 12%

↑ Indicates increase ↓ Indicates decrease ↔ Indicates no change or a mean increase or decrease of <10%

<sup>a</sup> Relative to steady-state administration of efavirenz (600 mg once daily for 9 days).

<sup>b</sup> SGC Soft Gelatin Capsule

<sup>c</sup> Not marketed in Canada

<sup>d</sup> Administered as ATRIPLA (efavirenz, emtricitabine and tenofovir DF fixed-dose combination)

## STORAGE AND STABILITY

SUSTIVA (efavirenz) capsules should be stored at 25°C; excursions permitted to 15 - 30°C [see USP Controlled Room Temperature.]

## **DOSAGE FORMS, COMPOSITION AND PACKAGING**

### **Dosage Forms**

SUSTIVA (efavirenz) is available as capsules for oral administration.

### **Composition and Packaging**

Capsules contain either 50 mg or 200 mg of efavirenz.

*Capsules 200 mg* are gold, reverse printed with “SUSTIVA” on the body and imprinted “200 mg” on the cap. Available in bottles containing 90 capsules.

*Capsules 50 mg* are gold and white, printed with “SUSTIVA” on the gold cap and purple oval reverse printed with “50 mg” on the white body. Available in bottles containing 30 capsules.


Capsules contain either 50 mg or 200 mg of efavirenz and the following inactive ingredients: sodium starch glycolate, lactose monohydrate, sodium lauryl sulfate and magnesium stearate. The capsule shell contains the following inactive ingredients and dyes: gelatin, sodium lauryl sulfate, titanium dioxide and/or yellow iron oxide. The capsule shells may also contain silicon dioxide. The capsules are printed with ink containing carmine, FD&C Blue No. 2 and titanium dioxide.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION

#### Drug Substance

Proper Name:	Efavirenz (USAN)
Chemical Name:	(S) -6- chloro-4-(cyclopropylethynyl)-1,4-dihydro-4-(trifluoromethyl)-2H-3,1-benzoxazin-2-one.
Molecular Formula:	C <sub>14</sub> H <sub>9</sub> ClF <sub>3</sub> NO <sub>2</sub>
Molecular Mass:	315.68
Structural Formula:	



#### Physicochemical Properties

Description:	Efavirenz is a white to slightly pink crystalline powder.
Solubility:	Efavirenz is practically insoluble in water at a concentration of <10 µg/mL.
Partition coefficient:	The (octanol/water) partition coefficient is determined to be P=5.4.
Melting point:	The melting point is 137.2 ± 1.4°C.
pKa:	The pKa is 10.2.

## CLINICAL TRIALS

### Clinical Studies

The principal efficacy analyses compared the durability of virologic suppression by assessing the proportion of subjects responding to treatment with HIV RNA less than the assay limit. Plasma HIV RNA levels were quantified using the AMPLICOR HIV-1 MONITOR assay. The standard assay with a limit of 500 copies/mL was used in study 364. In studies 006 and 020, standard assay with a limit of 400 copies/mL and ultrasensitive assay with a limit of 50 copies/mL were used. During study 006, version 1.5 of the AMPLICOR assay was introduced in Europe to enhance detection of non-clade B virus.

The secondary analyses compared the magnitude and durability of the change in plasma HIV-RNA levels and CD4 cell counts from baseline.

In the analysis, patients who terminated the study early for any reason or who had a missing HIV-RNA measurement that was either preceded or followed by a measurement above the limit of assay quantification were considered to have HIV-RNA above 400 copies/mL, or above 50 copies/mL in the case of the Ultrasensitive assay, at the missing time points.

***Study 006: SUSTIVA (efavirenz) + indinavir or SUSTIVA + zidovudine + lamivudine versus indinavir + zidovudine + lamivudine in antiretroviral-naïve or NRTI-experienced (lamivudine-naïve) patients:***

Study 006 is a randomized open label trial to evaluate the plasma HIV-RNA suppression achieved by SUSTIVA in combination with either indinavir (IDV) or with zidovudine (ZDV) + lamivudine (3TC) compared to indinavir plus zidovudine + lamivudine in HIV-infected patients naïve to lamivudine, protease inhibitors and NNRTIs. Study 006 was designed as an equivalency trial. Patients were randomized to one of three treatment regimens: SUSTIVA (600 mg daily) + indinavir (1000 mg q8h) or SUSTIVA (600 mg daily) + zidovudine (300 mg q12h) + lamivudine (150 mg q12h) versus indinavir (800 mg q8h) + zidovudine (300 mg q12h) + lamivudine (150 mg q12h). The 1266 patients enrolled in the study had a mean age of 36.5 years (range 18-81); 60% were caucasian and 83% were male. The mean baseline CD4 count was 341 cells/mm<sup>3</sup>, mean HIV RNA plasma level was 4.8 log<sub>10</sub> copies/mL. Forty-eight and 168 week data are presented in Table 11. Through 48 and 168 weeks of therapy, increases in CD4 cell counts were not significantly different between treatment arms.

**Table 11**  
**Study 006 - Summary of Key Efficacy Results - Week 48 and Week 168**

<b>Treatment Regimen</b>	<b>SUSTIVA + IDV</b>		<b>SUSTIVA + ZDV + 3TC</b>		<b>IDV + ZDV + 3TC</b>	
Total N Randomized	N = 429		N = 422		N = 415	
<b>Patients with Plasma HIV-RNA &lt;400 copies/mL - Amplicor Assay</b>						
Responder / Evaluable (%)						
	Week 48	Week 168	Week 48	Week 168	Week 48	Week 168
TLOVR <sup>a</sup>	246 / 429 (57)	171 / 429 (40)	293 / 422 (69)	203 / 422 (48)	206 / 415 (50)	123 / 415 (30)
TTF <sup>b</sup>	234 / 429 (55)	161 / 429 (38)	270 / 422 (64)	190 / 422 (45)	186 / 415 (45)	114 / 415 (27)
VR-OC	249 / 294 (85)	172 / 181 (95)	300 / 317 (95)	216 / 228 (95)	216 / 253 (85)	127 / 140 (91)
Difference Estimate (97.5% CI) at Week 168						
Treatment Regimen	SUSTIVA+IDV - IDV+ZDV+3TC			SUSTIVA+ZDV+3TC - IDV+ZDV+3TC		
TLOVR <sup>a</sup>	10.2 (2.9, 17.6) p = 0.0018 <sup>c</sup>			18.5 (10.9, 26.0) p < 0.0001 <sup>d</sup>		
TTF <sup>b</sup>	10.1 (2.8, 17.3) p = 0.0018 <sup>c</sup>			17.6 (10.1, 25.0) p < 0.0001 <sup>d</sup>		
VR-OC	4.3 (-2.1, 10.7) p = 0.1293 <sup>c</sup>			4.0 (-2.0, 10.1) p = 0.1365 <sup>d</sup>		
<b>Patients with Plasma HIV-RNA &lt;50 copies/mL - Ultrasensitive Assay</b>						
Responder / Evaluable (%)						
	Week 48	Week 168	Week 48	Week 168	Week 48	Week 168
TLOVR <sup>a</sup>	209 / 429 (49)	134 / 429 (31)	268 / 422 (64)	180 / 422 (43)	177 / 415 (43)	96 / 415 (23)
TTF <sup>b</sup>	204 / 429 (48)	128 / 429 (30)	255 / 422 (60)	169 / 422 (40)	166 / 415 (40)	89 / 415 (21)
VR-OC	214 / 294 (73)	158 / 181 (87)	273 / 317 (86)	207 / 228 (91)	189 / 253 (75)	113 / 140 (81)
Difference Estimate (97.5% CI) at Week 168						
Treatment Regimen	SUSTIVA+IDV - IDV+ZDV+3TC			SUSTIVA+ZDV+3TC - IDV+ZDV+3TC		
TLOVR <sup>a</sup>	8.1 (1.2, 15.0) p = 0.0082 <sup>c</sup>			19.5 (12.2, 26.8) p < 0.0001 <sup>d</sup>		
TTF <sup>b</sup>	8.4 (1.6, 15.1) p = 0.0053 <sup>c</sup>			18.6 (11.4, 25.8) p < 0.0001 <sup>d</sup>		
VR-OC	6.6 (-2.6, 15.7) p = 0.1070 <sup>c</sup>			10.1 (2.0, 18.2) p = 0.0053 <sup>d</sup>		
<b>Change from Baseline - HIV RNA level (log<sub>10</sub> c/mL)</b>						
	Week 48	Week 168	Week 48	Week 168	Week 48	Week 168
N	289	165	310	210	251	133
Mean (se)	-2.65 (0.055)	-2.98 (0.058)	-2.91 (0.041)	-3.07 (0.047)	-2.65 (0.059)	-2.88 (0.068)
Median	-2.84	-3.08	-2.94	-3.09	-2.82	-2.96
<b>Change from Baseline - CD4 Cell Counts (cells/mm<sup>3</sup>)</b>						
	Week 48	Week 168	Week 48	Week 168	Week 48	Week 168
N	256	158	279	205	228	129
Mean (se)	191 (9.4)	319 (14.9)	190 (10.1)	329 (14.4)	180 (11.6)	329 (21.1)
Median	170	300	179	292	157	292

TLOVR = Time to Loss of Virologic Response

A minimum of two consecutive HIV RNA measurements < LOQ maintained through end of study without intervening confirmed rebounds or treatment discontinuations. If the last measurement on-study was > LOQ, the

subject was considered a failure for that visit. Deaths, loss to follow-up, and changes in antiretroviral therapy also counted as failure in the TLOVR algorithm.

TTF = Time to Treatment Failure

Loss of HIV RNA suppression (confirmed RNA  $\geq$  400 copies/mL), development of a CDC Class C AIDS-defining event, treatment discontinuation or start of alternative HIV treatment after having two consecutive HIV RNA determinations  $<$  LOQ, or failure to virologically suppress or failure to receive study medication after randomization up to the reported visit week.

VR-OC = Virologic Response - Observed Cases

Classified subjects who remained on treatment according to a single HIV RNA measurement, either  $<$  LOQ or  $\geq$  LOQ closest to the scheduled visit and within a predefined visit window. Only those subjects who remained on treatment at the time of their visit week were included. Subjects with HIV RNA  $\geq$  LOQ were considered failures. Subjects who remained on treatment and were missing a measurement were classified as responders only if their immediately previous and subsequent viral loads met the VR-OC criteria for response.

- <sup>a</sup> Proportion responding using the TLOVR definition of response
- <sup>b</sup> Proportion responding using the TTF definition of response
- <sup>c</sup> Statistical difference between SUSTIVA + IDV and IDV + ZDV + 3TC, at Week 168
- <sup>d</sup> Statistical difference between SUSTIVA + ZDV + 3TC and IDV + ZDV + 3TC, at Week 168

HIV-RNA status and the reasons for discontinuations of randomized treatment are summarized (Table 12).

**Table 12**  
**Outcomes of Randomized Treatment Through 48 and 168 Weeks, Study 006**

Outcome	SUSTIVA + IDV n = 429		SUSTIVA + ZDV + 3TC n = 422		IDV + ZDV + 3TC n = 415	
	Week 48	Week 168	Week 48	Week 168	Week 48	Week 168
Responder <sup>a</sup>	57%	40%	69%	48%	50%	30%
Virologic failure <sup>b</sup>	15%	19%	6%	12%	13%	17%
Discontinued for adverse events	5%	8%	7%	8%	15%	20%
Discontinued for other reasons <sup>c</sup>	23%	32%	17%	31%	22%	32%
CD4 + cell count (cells/mm <sup>3</sup> )						
Observed subjects (n)	256	158	279	205	228	129
Mean change from baseline	191	319	190	329	180	329

- <sup>a</sup> Patients achieved and maintained confirmed HIV-1 RNA  $<$ 400 copies/mL through Week 48 or Week 168.
- <sup>b</sup> Includes patients who rebounded, patients who were on study at Week 48 and failed to achieve confirmed HIV-1 RNA  $<$ 400 copies/mL at time of discontinuation, and patients who discontinued due to lack of efficacy.
- <sup>c</sup> Includes consent withdrawn, lost to follow-up, noncompliance, never treated, missing data, protocol violation, death, and other reasons. Patients with HIV-1 RNA levels  $<$ 400 copies/mL who chose not to continue in the voluntary extension phases of the study were censored at date of last dose of study medication.

A Kaplan-Meier analysis of time to loss of virologic response (HIV RNA  $<$ 400 copies/mL) suggests that both the trends of virologic response and differences in response continue through 4 years.

**Study ACTG 364: SUSTIVA in combination with nelfinavir (NFV) in NRTI-experienced patients:**

ACTG 364 is a randomized, double-blind, placebo-controlled 48-week study in NRTI-experienced patients who had completed two prior ACTG studies. One-hundred and ninety-five HIV-infected patients (mean age 41 years [range 18-76], 74% Caucasian, 88% male) received NRTIs in combination with SUSTIVA (600 mg daily), or nelfinavir (750 mg TID), or SUSTIVA (600 mg daily) plus nelfinavir in a randomized double-blinded manner. Upon entry into the study, all patients were assigned a new open label NRTIS regimen, which was dependent on their previous NRTIS treatment experience. Through 48 weeks of therapy, there was no significant difference in the mean CD4 cell count between the treatment arms. Overall efficacy results are summarized in Table 13.

**Table 13**  
**ACTG 364- Summary of Key Efficacy Results -Week 48**

	NFV + 2NRTIs	SUSTIVA + 2NRTIs	NFV + SUSTIVA + 2NRTIs
Total N Randomized	66	65	65
<b>Patients with Plasma HIV-RNA &lt;500 copies/mL (95% CI) - Amplicor Assay</b>			
LOCF	23/66 (34.8%) (23.4, 46.3)	39/65 (60.0%) ‡ (48.1, 71.9)	49/65 (75.4%)* # † (64.9, 85.9)
NC=F	19/63 (30.2%) (18.8, 41.5)	36/62 (58.1%) (45.8, 70.3)	45/64 (70.3%)* ‡ † (59.1, 81.5)
<b>Mean Change from Baseline - Log10 Transformed Plasma HIV-RNA (SEM)</b>			
Amplicor (LOCF)	-0.45 (0.09) <sup>†</sup> N = 66	-0.72 (0.09) <sup>†</sup> N = 65	-0.87 (0.10) <sup>†</sup> N = 62
<b>Mean Change from Baseline - CD4 Counts (SEM)</b>			
LOCF	93.8 (13.6) <sup>†</sup> N = 66	113.8 (21.0) <sup>†</sup> N = 65	107.4 (17.9) <sup>†</sup> N = 63

\* Statistically significant difference between NFV+2NRTIs and NFV+SUSTIVA+2NRTIs (p≤0.05)

# Statistically significant difference between SUSTIVA+2NRTIs and NFV+SUSTIVA+2NRTIs (p≤0.05)

† Statistically significant change from baseline (p≤0.05)

‡ Statistically significant difference between NFV+2 NRTIs and SUSTIVA+2NRTIs (p≤0.05).

† † Statistically significant difference among treatment groups.

LOCF: This analysis is based on the last obtained plasma HIV-RNA or CD4 cell count measurement for each patient randomized into the study (Intent-to-Treat: Last Observation Carried Forward analysis). NC=F: Noncompleter = Failure. In this analysis, any patient with viral load above quantifiable levels using a specified assay, or whose viral load data are missing at some time point of the analysis, is considered to be a virologic failure (i.e., HIV-RNA ≥500 copies/mL using the Roche Amplicor™ assay), unless the patient's viral load measurements both before and after the missing data point are below the assay limit of quantification.

CI: Confidence Interval

HIV-RNA status and the reasons for discontinuations of randomized treatment are summarized (Table 14).

**Table 14**  
**Study ACTG 364 - Outcomes of Randomized Treatment Through 48 Weeks**

<b>Outcome</b>	<b>SUSTIVA + NFV + NRTIs N=65</b>	<b>SUSTIVA + NRTIs N=65</b>	<b>NFV + NRTIs N=66</b>
HIV-RNA <500 copies/mL <sup>1</sup>	70%	58%	30%
HIV-RNA ≥500 copies/mL <sup>2,3</sup>	13%	27%	60%
Discontinuations for Adverse Events <sup>3</sup>	3%	3%	5%
Discontinuations for Other Reasons <sup>3</sup>	9%	2%	0%
On Treatment with Missing HIV-RNA Value <sup>3</sup>	5%	10%	5%
<b>Total</b>	<b>100%</b>	<b>100%</b>	<b>100%</b>

<sup>1</sup> Corresponds to rates at 48 weeks in Table 13

<sup>2</sup> Includes discontinuation due to virologic failure at or before 48 weeks

<sup>3</sup> Treatment Failure in the Analysis

***Study 020: Protease Inhibitor + Two NRTIs with/without SUSTIVA in NRTI-experienced patients:***

Study 020 was a randomized, double-blind, placebo-controlled 24-week study in NRTI-experienced, protease inhibitor and NNRTI-naive patients designed to compare quadruple therapy consisting of SUSTIVA + indinavir + 2 nucleoside analogue reverse transcriptase inhibitors versus triple therapy consisting of indinavir + 2 NRTIs at 24 weeks of treatment. Patients were randomized to receive either SUSTIVA (600 mg daily) + indinavir (1000 mg q8h) + 2 NRTIs or indinavir (800 mg q8h) + 2 NRTIs. Sixty-seven percent of the 327 patients (mean age 38.5 years [range 20-69], 52% Caucasian, 83% male) changed their NRTI regimen at study initiation. Mean baseline CD4 count was 328 cells/mm<sup>3</sup>, and mean HIV-RNA plasma level was 4.41 log<sub>10</sub> copies/mL. Through 24 weeks of therapy, there was no significant difference in the mean CD4 cell count between the treatment arms. Mean increases in CD4 cell count at 24 weeks were 104 cells/mm<sup>3</sup> for patients treated with SUSTIVA+IDV+NRTIs, and 77 cells/mm<sup>3</sup> for patients treated with IDV+NRTIs. The Noncompleter=Failure analysis at 24 weeks showed no difference between the treatment groups using a cutoff of 400 copies/mL and a significant difference between treatment groups using a cutoff of 50 copies/mL. Efficacy results are summarized in Table 15.

**Table 15**  
**Study 020 - Summary of Key Efficacy Results - Week 24**

	<b>SUSTIVA + NRTIs</b>	<b>IDV + NRTIs</b>
Total N Randomized	157	170
<b>Patients with Plasma HIV-RNA &lt;400 copies/mL (95% CI) - Amplicor Assay</b>		
LOCF	107/157 (68.2%)* (60.5, 75.8)	89/170 (52.4%) (44.6, 60.2)
NC=F	93/156 (59.6%) (51.6, 67.6)	86/169 (50.9%) (43.1, 58.7)
Observed	93/112 (83.0%)* (75.6, 90.4)	86/132 (65.2%) (56.6, 73.7)
<b>Patients with Plasma HIV-RNA &lt;50 copies/mL (95% CI) - Ultrasensitive Assay</b>		
LOCF	79/156 (50.6%)* (42.5, 58.8)	65/168 (38.7%) (31.0, 46.4)
NC+F	77/156 (49.4%)* (41.2, 57.5)	63/168 (37.5%) (29.9, 45.1)
Observed	77/112 (68.8%)* (59.7, 77.8)	63/132 (47.7%) (38.8, 56.6)
<b>Mean Change from Baseline - Log<sub>10</sub> Transformed Plasma HIV-RNA (SEM)</b>		
Amplicor (LOCF)	-1.45 (0.08)* † N = 147	-1.12 (0.08)† N = 158
Ultrasensitive** (LOCF)	2.25 (0.10)* † N = 120	-1.72 (0.11)†
<b>Mean Change from Baseline - CD4 Counts (SEM)</b>		
LOCF	104.4 (9.1)* † N = 151	76.9 (9.9)† N = 158

\* Statistically significant difference between treatments (p≤0.05)

† Statistically significant change from baseline (p≤0.05)

\*\* Ultrasensitive Assay is an unvalidated experimental method.

LOCF: This analysis is based on the last obtained plasma HIV-RNA or CD4 cell count measurement for each patient randomized into the study (Intent-to-Treat: Last Observation Carried Forward analysis).

NC=F: Noncompleter = Failure. In this analysis, any patient with viral load above quantifiable levels using a specified assay, or whose viral load data are missing at some time point of the analysis, is considered to be a virologic failure (i.e., HIV-RNA ≥400 copies/mL using the Roche Amplicor™ assay), unless the patient's viral load measurements both before and after the missing data point are below the assay limit of quantification.

Observed: Analysis is based on all data available at the specified time point (on-treatment analysis). Missing data is not accounted for in this analysis.

CI: Confidence Interval

HIV-RNA status and the reasons for discontinuations of randomized treatment are summarized (Table 16).

**Table 16**  
**Study 020 - Outcomes of Randomized Treatment Through 24 Weeks**

<b>Outcome</b>	<b>SUSTIVA + IDV + NRTIs N = 157</b>	<b>IDV + NRTIs N = 170</b>
HIV-RNA <400 copies/mL [<50 copies/mL]	60% [51%]	51% [39%]
HIV-RNA ≥400 copies/mL [≥50 copies/mL] <sup>1,2</sup>	11% [20%]	26% [38%]
Discontinuations for Adverse Events <sup>2</sup>	11%	5%
Discontinuations for Other Reasons <sup>2</sup>	18%	16%
On Treatment with Missing HIV-RNA Value <sup>2</sup>	0%	2%
<b>Total</b>	<b>100%</b>	<b>100%</b>

<sup>1</sup> Includes discontinuation due to virologic failure at or before 48 weeks

<sup>2</sup> Treatment Failure in the Analysis

## DETAILED PHARMACOLOGY

### Pharmacokinetics

#### Absorption

Peak efavirenz plasma concentrations of 1.6-9.1 µM were attained by 5 hours following single oral doses of 100 mg to 1600 mg administered to uninfected volunteers. Dose-related increases in C<sub>max</sub> and AUC were seen for doses up to 1600 mg; the increases were less than proportional suggesting diminished absorption at higher doses. Time to peak plasma concentrations (3-5 hours) did not change following multiple dosing and steady-state plasma concentrations were reached in 6-10 days.

In HIV-infected patients at steady-state, mean C<sub>max</sub>, mean C<sub>min</sub>, and mean AUC were dose proportional following 200 mg, 400 mg, and 600 mg daily doses. In 35 patients receiving SUSTIVA 600 mg daily, mean steady-state C<sub>max</sub> was 12.9 µM, mean steady state C<sub>min</sub> was 5.6 µM, and mean AUC was 184 µM/h.

#### Effect of Food on Oral Absorption

Administration of a single 600 mg dose of efavirenz capsules with a high fat/high caloric meal (894 kcal, 54 g fat, 54% calories from fat) or a reduced fat/normal caloric meal (440 kcal, 2 g fat, 4% calories from fat) was associated with a mean increase of 22% and 17% in efavirenz AUC and a mean increase of 39% and 51% in efavirenz C<sub>max</sub>, respectively, relative to the exposures achieved when given under fasted conditions. (See [DOSAGE AND ADMINISTRATION](#).)

#### Distribution

Efavirenz is highly bound (approximately 99.5-99.75%) to human plasma proteins, predominantly albumin. In HIV-1 infected patients (N=9) who received SUSTIVA (efavirenz) 200 to 600 mg once daily for at least one month, cerebrospinal fluid concentrations ranged from

0.26 to 1.19% (mean 0.69%) of the corresponding plasma concentration. This proportion is approximately 3-fold higher than the non-protein-bound (free) fraction of efavirenz in plasma.

### Metabolism

Studies in humans and *in vitro* studies using human liver microsomes have demonstrated that efavirenz is principally metabolized by the cytochrome P450 system to hydroxylated metabolites with subsequent glucuronidation of these hydroxylated metabolites. These metabolites are essentially inactive against HIV-1. The *in vitro* studies suggest that CYP3A4 and CYP2B6 are the major isozymes responsible for efavirenz metabolism. *In vitro* studies have shown that efavirenz inhibited P450 isozymes 2C9, 2C19, and 3A4 with  $K_i$  values (8.5-17.0  $\mu\text{M}$ ) in the range of observed efavirenz plasma concentrations. In *in vitro* studies efavirenz did not inhibit CYP2E1 and inhibited CYP2D6 and CYP1A2 ( $K_i$  values 82-160  $\mu\text{M}$ ) only at concentrations well above those achieved clinically.

Efavirenz plasma exposure may be increased in patients with the homozygous G516T genetic variant of the CYP2B6 isozyme. The clinical implications of such an association are unknown; however, the potential for an increased frequency and severity of efavirenz-associated adverse events cannot be excluded.

Efavirenz has been shown to induce P450 enzymes, resulting in the induction of its own metabolism. Multiple doses of 200-400 mg per day for 10 days resulted in a lower than predicted extent of accumulation (22-42% lower) and a shorter terminal half-life of 40-55 hours (single dose half-life 52-76 hours). The degree of CYP3A4 induction is expected to be similar between a 400 mg and 600 mg dose of efavirenz based on pharmacokinetic interaction studies in which daily 400 mg or 600 mg efavirenz doses in combination with indinavir did not appear to cause any further reduction of indinavir AUC compared to a 200 mg dose of efavirenz.

### Elimination

Efavirenz has a long terminal half-life of 52-76 hours after single doses and 40-55 hours after multiple doses. A one-month mass balance/excretion study was conducted using 400 mg per day with a  $^{14}\text{C}$ -labelled dose administered on Day 8. Approximately 14-34% of the radiolabel was recovered in the urine and 16-61% was recovered in the feces. Nearly all of the urinary excretion of the radiolabelled drug was in the form of metabolites. Efavirenz accounted for the majority of the total radioactivity measured in feces.

### Demographics

The pharmacokinetics of efavirenz in patients appear to be similar between men and women and among the ethnic groups studied.

### Hepatic Enzyme Induction

Efavirenz has been shown *in vivo* to cause hepatic enzyme induction, thus increasing the biotransformation of some drugs metabolized by CYP3A4 and CYP2B6. *In vitro* studies have demonstrated that efavirenz inhibits 2C9, 2C19 and 3A4 isozymes with  $K_i$  values (8.5-17  $\mu\text{M}$ ) in the range of observed efavirenz plasma concentrations. In *in vitro* studies, efavirenz did not

inhibit CYP2E1 and inhibited CYP2D6 and CYP1A2 ( $K_i$  values 82-160  $\mu\text{M}$ ) only at concentrations well above those achieved clinically. The effects on CYP3A4 activity are expected to be similar between 200 mg, 400 mg and 600 mg doses of efavirenz. Coadministration of efavirenz with drugs primarily metabolized by 2C9, 2C19 and 3A4 isozymes may result in altered plasma concentrations of the coadministered drug. Drugs which induce CYP3A4 activity would be expected to increase the clearance of efavirenz resulting in lowered plasma concentrations.

## VIROLOGY

Efavirenz is a selective non-nucleoside reverse transcriptase inhibitor (NNRTI) of human immunodeficiency virus type 1 (HIV-1). Efavirenz is predominantly a non-competitive inhibitor of HIV-1 reverse transcriptase (RT). HIV-2 RT and human cellular DNA polymerases  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$  are not inhibited by concentrations of efavirenz well in excess of those achieved clinically.

***In vitro* HIV Susceptibility:** The clinical significance of *in vitro* susceptibility of HIV-1 to efavirenz has not been established. The *in vitro* antiviral activity of efavirenz was assessed in lymphoblastoid cell lines, peripheral blood mononuclear cells (PBMCs) and macrophage/monocyte cultures enriched from PBMCs. The 90-95% inhibitory concentration ( $\text{IC}_{90-95}$ ) of efavirenz for wild type laboratory adapted strains and clinical isolates ranged from 1.7 to  $\leq 25\text{nM}$ . EFV demonstrated antiviral activity against most non-clade B isolates (subtypes A, AE, AG, C, D, F, G, J, N), but had reduced antiviral activity against group O viruses. EFV demonstrated additive antiviral activity without cytotoxicity against HIV-1 in cell culture when combined with the NNRTIs delavirdine (DLV) and nevirapine (NVP), NRTIs (abacavir, didanosine, emtricitabine, lamivudine [LAM], stavudine, tenofovir, zalcitabine, zidovudine [ZDV]), PIs (amprenavir, indinavir [IDV], lopinavir, nelfinavir, ritonavir, saquinavir), and the fusion inhibitor enfuvirtide. EFV demonstrated additive to antagonistic antiviral activity *in vitro* with atazanavir. EFV was not antagonistic with adefovir, used for the treatment of hepatitis B virus infection, or ribavirin, used in combination with interferon for the treatment of hepatitis C virus infection.

**Resistance:** HIV-1 isolates with reduced susceptibility to efavirenz ( $>380$ -fold increase in  $\text{IC}_{90}$ ) compared to baseline can emerge rapidly in cell culture in the presence of drug. Genotypic characterization of these viruses identified mutations resulting in single amino acid substitutions L100I or V179D, double substitutions L100I/V108I, and triple substitutions L100I/V179D/Y181C in RT. Phenotypic ( $N=26$ ) changes in evaluable HIV-1 isolates and genotypic ( $N=104$ ) changes in plasma virus from selected patients treated with efavirenz in combination with IDV, or with ZDV plus lamivudine were monitored. Clinical isolates with reduced susceptibility in cell culture to efavirenz have been obtained. One or more RT mutations at amino acid positions 98, 100, 101, 103, 106, 108, 188, 190 and 225, and 227 were observed in all 102 of 104 patients with a frequency of at least 9% compared to baseline. The mutation at RT amino acid position 103 (lysine to asparagine) was the most frequently observed ( $\geq 90\%$ ). A mean loss in susceptibility ( $\text{IC}_{90}$ ) to efavirenz of 47-fold was observed in 26 clinical isolates. Five clinical

isolates were evaluated for both genotypic and phenotypic changes from baseline. Decreases in efavirenz susceptibility (range from 9 to >312-fold increase in IC<sub>90</sub>) were observed for these isolates in cell culture compared to baseline. All 5 isolates possessed at least one of the efavirenz-associated RT mutations. The clinical relevance of phenotypic and genotypic changes associated with efavirenz therapy has not been established. Long-term resistance surveillance (average 52 weeks, range 4-106 weeks) analyzed 28 matching baseline and virologic failure isolates. Sixty-one percent (17/28) of these failure isolates had decreased EFV susceptibility in cell culture with a median 88-fold change in EFV susceptibility (IC<sub>50</sub> value) from reference. The most frequent NNRTI mutation to develop in these patient isolates was K103N (54%). Other NNRTI mutations that developed included L100I (7%), K101E/Q/R (14%), V108I (11%), G190S/T/A (7%), P225H (18%), and M230I/L (11%).

**Cross-Resistance:** Rapid emergence of HIV-1 strains that are cross-resistant to non-nucleoside RT inhibitors has been observed in cell culture. Thirteen clinical isolates previously characterized as efavirenz-resistant were also phenotypically resistant to nevirapine and delavirdine in cell culture compared to baseline. DLV- and/or NVP-resistant clinical viral isolates with NNRTI resistance-associated substitutions (A98G, L100I, K101E/P, K103N/S, V106A, Y181X, Y188X, G190X, P225H, F227L, or M230L) showed reduced susceptibility to EFV in cell culture. Clinically derived ZDV-resistant HIV-1 isolates tested in cell culture retained susceptibility to efavirenz. Cross-resistance between efavirenz and HIV protease inhibitors is unlikely because of the different enzyme targets involved.

## TOXICOLOGY

### Acute Toxicity

The oral Minimum Lethal Dose (MLD) of efavirenz in female rats ranged from 250 mg/kg to 500 mg/kg, whereas the oral MLD of efavirenz in male rats was 1000 mg/kg. The most prominent clinical signs attributed to efavirenz treatment in rats were ataxia and decreased motor activity, and were observed, in general, at doses  $\geq$ 250 mg/kg. The MLD in female and male rats given intraperitoneal injections of efavirenz was 250 mg/kg and 500 mg/kg, respectively. The MLD in mice given intraperitoneal injections of efavirenz was similar to that in rats (250 mg/kg in both male and female mice).

### **Efavirenz Toxicokinetics in Preclinical Studies**

Plasma efavirenz concentrations achieved at maximally tolerated doses in rats were lower than those achieved in humans given efavirenz (in rats given 250 mg/kg bid of efavirenz the AUC was 38  $\mu\text{M}\cdot\text{h}$  in males and 84  $\mu\text{M}\cdot\text{h}$  in females). This was due, in part, to the autoinduction caused by efavirenz in rats. The resulting plasma half-life of efavirenz in rats is  $\approx$  0.8 to 1.9 hours, compared to a half-life of >40 hours in humans. Due to the low plasma efavirenz concentrations attained in rats, rats were a poor model in which to assess the toxicity of the parent drug. However, plasma concentrations of the primary efavirenz metabolite (the 8-OH-glucuronide conjugate) were 50- to 75-fold higher in rats given 250 mg/kg bid than in humans

given therapeutic doses of efavirenz (the 8-OH-glucuronide conjugate was present at millimolar concentrations in the plasma of rats).

Chronic toxicity studies were conducted in cynomolgus monkeys in which the plasma efavirenz AUC values were higher than those attained in humans. AUC values achieved in cynomolgus monkeys given 45 mg/kg bid and 75 mg/kg bid during subchronic and chronic studies were  $\approx$  1.5- and 5-fold higher, respectively, than those achieved in humans given 600 mg/day (AUC values of 283  $\mu\text{M}\cdot\text{h}$  and 907  $\mu\text{M}\cdot\text{h}$ , respectively). In addition, the metabolic disposition of efavirenz in cynomolgus monkeys is similar to that in humans, although efavirenz is more extensively metabolized in cynomolgus monkeys than in humans.

### **Long-term Toxicity**

The toxicity and toxicokinetics of efavirenz were evaluated in rats (duration of study  $\leq$ 6 months), in rhesus monkeys (duration of study 1 month) and in cynomolgus monkeys (duration of study  $\leq$ 1 year). In addition, a 5 week oral toxicity study was conducted in newborn rhesus monkeys. From the subacute/chronic toxicological studies the following adverse effects have been noted following treatment with efavirenz.

### **Nephrotoxicity**

The most prominent toxicologic finding in rats given efavirenz was nephrotoxicity. Rats treated with  $\geq$ 250 mg/kg bid efavirenz in the 3-month oral toxicity study showed mild to moderate renal cortical epithelial cell necrosis, associated with intraluminal casts, proteinaceous debris, and tubular dilatation. In addition, late-stage renal changes (occasionally leading to renal failure) occurred secondary to tubular obstruction and were characterized by cystic tubular dilatation and degeneration. Treatment of rats with doses  $\geq$ 500 mg/kg bid occasionally resulted in death due to acute tubular necrosis. Renal lesions elicited by 1 month of dosing with 250 mg/kg bid of efavirenz were shown to be fully reversible following 1-month recovery period. The incidence of and severity of renal lesions was greater in male than female rats given equivalent efavirenz doses. The no effect dose for renal toxicity after  $\geq$ 3 months of dosing was 100 mg/kg bid in female rats and 30 mg/kg bid in male rats.

Nephrotoxicity did not occur in cynomolgus monkeys given 75 mg/kg bid (AUC=907  $\mu\text{M}\cdot\text{h}$ ) for 3 months to 1 year or in rhesus monkeys given 100 mg/kg/day for 1 month (AUC=212  $\mu\text{M}\cdot\text{h}$ ). Average plasma efavirenz concentrations in monkeys in these studies exceeded those in rats given nephrotoxic doses of efavirenz.

Special toxicity studies implicated the formation of an efavirenz glutathione conjugate in the nephrotoxicity caused by efavirenz in rats. In these studies, interventions to decrease the formation and/or the catabolic products of glutathione conjugates resulted in decreased efavirenz-induced nephrotoxicity. Glutathione conjugates of efavirenz are not found in cynomolgus monkeys or in humans given efavirenz. Therefore, the nephrotoxicity elicited by efavirenz in rats is considered to be species specific.

### **Changes in the Biliary System**

- a) **Biliary Fibrosis in Rats:** An increased incidence of biliary fibrosis occurred in the liver of rats after oral gavage or dietary administration of doses  $\geq 500$  mg/kg/day. The severity of the change was minimal to mild and was sparsely distributed involving relatively few isolated bile ducts. In rats given efavirenz in the diet for 3 months the lesions were sometimes accompanied by multifocal biliary hyperplasia (a possible consequence of increased biliary pressure in the drainage areas where fibrosis had occurred). Biliary lesions were not observed in rats given 100 mg/kg bid by oral gavage for 6 months or 100 mg/kg/day in the diet for 3 months.
- b) **Biliary Hyperplasia in Cynomolgus Monkeys:** Minimal biliary hyperplasia (increase in the number of small caliber bile ducts) was observed in the liver of two of four male and two of four female monkeys in the high dose (75 mg/kg bid; AUC=907  $\mu\text{M}(\text{h})$ ) cynomolgus monkeys given efavirenz for 1 year. (Biliary hyperplasia was not seen in monkeys given this dose for 6 months or less.) There was no serum biochemical or histologic evidence of cholestasis and the change was not accompanied by fibrosis or histologic evidence of adjacent hepatocellular injury. In monkeys given the dose of 75 mg/kg bid, the mean plasma efavirenz AUC was approximately 5-fold greater than the AUC in humans given 600 mg/day of efavirenz (AUC=186  $\mu\text{M}(\text{h})$ ). The no effect level for biliary hyperplasia (45 mg/kg bid for  $\leq 1$  year efavirenz (AUC=283  $\mu\text{M}(\text{h})$ ) was approximately 1.5-fold greater than in humans given 600 mg/day of efavirenz.

In a subsequent 2 year study total daily doses of  $\geq 60$  mg/kg/day resulted in minimal to moderate biliary hyperplasia in almost all of the animals. Plasma efavirenz exposure levels (120 mg/kg/day, AUC 1871  $\mu\text{M}(\text{h})$  at 99 weeks) were higher in this study than in the 1 year study (see [table 17](#)).

**Table 17**

	Total Daily Dose	Cmax range	AUC range
1 year study (T95-10-4)	150 mg/kg/day	32 - 72 $\mu\text{M}$	489 - 1262 $\mu\text{M}\cdot\text{h}$
2 year study (T97-11-1)	150 mg/kg/day	42 - 159 $\mu\text{M}$	745 - 3173 $\mu\text{M}\cdot\text{h}$

The biliary hyperplasia was not associated with any fibrotic or degenerative changes. The cause of biliary hyperplasia in cynomolgus monkeys following treatment with efavirenz remains unknown.

### **Minimal Thyroid Follicular Cell Hypertrophy**

Minimal thyroid follicular hypertrophy was observed in cynomolgus monkeys given  $\geq 45$  mg/kg bid (AUC=283  $\mu\text{M}(\text{h})$ ) for 1 year but not in monkeys given 15 mg/kg bid (AUC=65  $\mu\text{M}(\text{h})$ ) for 1 year, or 100 mg/kg bid of efavirenz for  $\leq 6$  months or in rats. In a 2-year study in cynomolgus monkeys minimal-to-slight hypertrophy of thyroid follicular cells, was observed in the thyroid lobes of 1/11, 4/10, 8/10, and 1/5 monkeys given 0, 60, 150, or 150/80 mg/kg/day of DMP 266, respectively. In this study, thyroid follicular cell hypertrophy was not observed at the end of the 26-week recovery period. An increased clearance of  $^{125}\text{I}$ -thyroxine and elevations in serum thyroid stimulating hormone (TSH) were observed in cynomolgus monkeys given 75 mg/kg bid

of efavirenz for 1 month. The thyroid follicular cell hypertrophy may be due to the induction of UDP-glucuronyl transferase in this species (a rate limiting Phase II enzyme involved in the clearance of thyroxine).

### **ALT Elevations**

Slight increases in serum ALT but not AST activity were observed in individual cynomolgus monkeys given  $\geq 45$  mg/kg bid (AUC  $\geq 283$   $\mu\text{M}(\text{h})$ ). The greatest individual ALT elevation was approximately 3-fold above the highest concurrent control, with the majority of the ALT elevations being mild (no more than approximately 1.5-fold above the highest concurrent control value). There was no evidence of hepatocellular injury after a year of dosing. No drug-related elevations of ALT were noted in rats or rhesus monkeys treated with efavirenz.

### **APTT and PT Prolongations**

Slight increases in prothrombin time (PT) and activated partial thromboplastin time (APTT) were observed in some male rats given  $\geq 50$  mg/kg bid for 6 months. These prolongations were not associated with any gross or microscopic evidence of bleeding. No drug-related changes were found in the PT and APTT of female rats, nor were any prolongations observed in PT or APTT of male or female rats in studies of  $\leq 3$  months duration. The cause of these prolongations in male rats is unknown.

In cynomolgus monkeys, slightly prolonged APTT (up to approximately 10 seconds above the upper limit of the reference range) were observed in some monkeys given 45 mg/kg bid (AUC=283  $\mu\text{M}(\text{h})$ ) or 75 mg/kg bid (AUC=907  $\mu\text{M}(\text{h})$ ) of efavirenz for  $> 6$  months. The incidence and magnitude of the APTT prolongations increased with dose and the prolongations remained relatively constant over the course of the studies. Further investigation revealed a slight decrease in the activities of Factor XII in affected monkeys, and a slight decrease in the activity of Factor XI in the monkeys with the longest APTT. The decreased factor activity was not attributed to the presence of drug-induced inhibitor. Apart from decreased coagulation factor activities, no alteration in coagulation parameters, fibrinogen concentration, prothrombin time, or platelet count were noted. There was no evidence of gross or microscopic bleeding upon postmortem examination. The cause of decreased activities of Factor XII and/or XI is unknown.

### **Mutagenicity**

Efavirenz was negative in a battery of in vitro and in vivo genotoxicity assays. This included assays in four in vitro assay systems: 1) bacterial mutation assays in *Salmonella typhimurium* and *Escherichia coli*, 2) a Chinese hamster ovary (CHO) cell/hypoxanthine-guanine phosphoribosyl-transferase (HGPRT) forward mutation assays, 3) a chromosome aberration assay in human peripheral lymphocytes, and 4) a chromosome aberration assay in CHO cells, and in one in vivo system (mouse micronucleus assay). All assays were conducted employing maximally soluble or minimally toxic doses/concentrations of efavirenz.

### **Reproduction and Teratology**

Malformations were observed in 3 of 20 fetuses/infants from efavirenz-treated cynomolgus monkeys (versus 0 of 20 concomitant controls) in an ongoing developmental toxicity study. The pregnant monkeys were dosed throughout pregnancy (post-coital days 20-150) with efavirenz 60 mg/kg daily, a dose which resulted in plasma drug concentrations similar to those in humans given 600 mg/day. Anencephaly and unilateral anophthalmia were observed in one fetus, microphthalmia was observed in another fetus, and cleft palate was observed in a third fetus.

No malformations were observed in fetuses from efavirenz-treated rats; however, an increase in fetal resorptions and a slight increase in pup mortality was observed in rats at doses that produced peak plasma concentrations and AUC values in pregnant female rats similar to or lower than those achieved in humans at the recommended clinical dose. Efavirenz was not teratogenic or embryotoxic when given to pregnant rabbits.

#### Peri/Postnatal Toxicity

A 5-8% decrease in mean rat pup weights versus control mean rat pup weights, and a slight increase in rat pup mortality was observed at doses of 50 and 100 mg/kg bid in a study in which efavirenz was given to pregnant rats during gestation and through lactation until weaning (the 100 mg/kg bid dose produced peak plasma concentrations and AUC values in pregnant female rats similar to or lower than those achieved in humans at the recommended clinical dose). No efavirenz-related effects were observed on the fertility, mating behavior, sexual maturation, learning, or behavior of the F1 generation derived from female rats given 100 mg/kg bid.

#### Maternal/Fetal Efavirenz Exposure

Fetal exposure to efavirenz was documented in pregnant rats, rabbits and cynomolgus monkeys. Maternal and fetal blood concentrations were equivalent in pregnant rabbits and cynomolgus monkeys and fetal blood concentrations were approximately 25% to 49% lower than the corresponding maternal concentrations in pregnant rats. Results of these studies indicated that efavirenz crossed the placenta in all species tested.

#### Efavirenz Concentration in Milk

The excretion of efavirenz into rat milk was demonstrated. Efavirenz milk concentrations in rats were approximately 8-fold higher than corresponding maternal efavirenz plasma concentrations.

#### Male/Female Fertility Assessment

No efavirenz-related effects were observed on the fertility or reproductive performance of female rats given 100 mg/kg bid, or on the reproductive performance or sperm motility and morphology of male rats given 200 mg/kg bid.

#### Assessment of Toxicity in Infant and Neonatal Non-Human Primates

In a five-week oral infant rhesus toxicity study, (dosing initiated on Day 2 of life at 30 and 45 mg/kg bid), infant rhesus monkeys given 30 mg/kg bid exhibited a slight, transitory decrease in body weight gain in females and slight decreases in food intake in females. Doses of 45 mg/kg bid produced adverse clinical signs in infant rhesus monkeys (vomiting, lethargy, dehydration, poor appetite, and/or weakness) and slight decreases in the average amount of body weight gain.

No efavirenz-related hematology, serum biochemical, or histologic changes occurred at either dose.

### **Carcinogenesis**

In a 2-year carcinogenicity study, mice were given daily oral dosages of 25, 75, 150 or 300 mg/kg/day of efavirenz. Because efavirenz is rapidly cleared in mice, plasma drug exposure (as measured by AUC) at dosages  $\leq 150$  mg/kg/day was lower than that in humans given 600 mg/day of efavirenz. In mice given 300 mg/kg/day of efavirenz, plasma drug exposure (AUC) was approximately 1.7-fold the AUC in humans given 600 mg/day of efavirenz. In female mice, a statistically significant, dose-related increase in the incidence of hepatic tumors occurred at dosages  $\geq 75$  mg/kg/day and a statistically significant, non dose-related increase in pulmonary tumors occurred at dosages  $\geq 25$  mg/kg/day of efavirenz. Efavirenz did not increase the incidence of any tumor type in male mice. Given the lack of genotoxic activity of efavirenz, the relevance to humans of hepatocellular tumors in efavirenz-treated mice is not known.

In a 2-year carcinogenicity study, rats were given daily oral dosages of 25, 50 or 100 mg/kg/day of efavirenz. Plasma drug exposures (as measured by AUC) in rats given all dosages of efavirenz were substantially below those achieved in humans given 600 mg/day of efavirenz, and therefore may not reflect the carcinogenic potential of efavirenz in humans. The low plasma drug exposures attained in rats are a consequence of the extremely rapid metabolic clearance of efavirenz in this species. However, virtually all of the efavirenz metabolites formed in rats are also formed in humans and the levels of these metabolites attained in the rat carcinogenicity study were likely substantially higher than those achieved in humans. Therefore, the results of this carcinogenicity study do provide meaningful information on the potential carcinogenicity of these efavirenz metabolites even at relatively low multiples of the parent drug exposure. Efavirenz did not increase the incidence of any tumor type in rats.

Long-term carcinogenicity studies in mice and rats were carried out with efavirenz. Mice were dosed with 0, 25, 75, 150, or 300 mg/kg/day for 2 years. Incidences of hepatocellular adenomas and carcinomas and pulmonary alveolar/bronchiolar adenomas were increased above background in females. No increases in tumor incidence above background were seen in males. In studies in which rats were administered efavirenz at doses of 0, 25, 50, or 100 mg/kg/day for 2 years, no increases in tumor incidence above background were observed. The systemic exposure (based on AUCs) in mice was approximately 1.7-fold that in human receiving the 600 mg/day dose. The exposure in rats was lower than that in humans.

The findings from the completed efavirenz mouse carcinogenicity study may not represent a significant risk to patients based upon the following: An increase in the incidence of hepatic tumors in efavirenz-treated mice was not unexpected as efavirenz is known to induce hepatic drug-metabolizing enzyme activity and enzyme inducers are known to increase the incidence of hepatic tumors in rodents, but not in humans. While the cause of the increased incidence of pulmonary tumors is not known, this finding also may not constitute a significant risk for patients given efavirenz because: (1) efavirenz is not genotoxic, (2) the strain of mice used in these studies is documented to have a high spontaneous background incidence of this tumor type,

and (3) a decrease in the incidence of pulmonary tumors was observed in efavirenz-treated male mice. In male mice, plasma efavirenz concentrations were equal to or greater than in female mice.

## PART III: CONSUMER INFORMATION

Pr SUSTIVA®  
(efavirenz, USP)

This leaflet is Part III of a three-part “Product Monograph” published when SUSTIVA was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about SUSTIVA. Contact your doctor or pharmacist if you have any questions about the drug.

### ABOUT THIS MEDICATION

#### What is SUSTIVA?

- SUSTIVA is the brand name for the active ingredient efavirenz.
- SUSTIVA belongs to a class of anti-HIV medicines known as "non-nucleoside reverse transcriptase inhibitors" (NNRTIs or non-nukes).

#### What the medication is used for:

- Your doctor has prescribed SUSTIVA for you because you have been infected with HIV. SUSTIVA must always be taken in combination with other anti-HIV medicines (frequently referred to as "combination therapy").
- When taken with other anti-HIV medicines, SUSTIVA has been shown to reduce viral load and increase the number of CD4 cells (a type of immune cell in blood). SUSTIVA may not have these effects in every patient.

#### Does SUSTIVA cure HIV or AIDS?

- SUSTIVA is not a cure for HIV nor Acquired Immunodeficiency Syndrome (AIDS). People taking SUSTIVA may still develop infections or other illnesses associated with HIV.
- It is very important that you remain under the constant care of your doctor while taking SUSTIVA.

#### Does SUSTIVA reduce the risk of passing HIV to others?

- SUSTIVA has not been shown to reduce the risk of passing HIV to others through sexual contact or blood contamination. It is important to continue to practice safe sex and not use or share dirty needles.

#### What it does:

- SUSTIVA fights Human Immunodeficiency Virus (HIV) infection by reducing the amount of virus in the blood (called "viral load").

#### When it should not be used:

- Do not take SUSTIVA if you know you are allergic to any of the ingredients in the SUSTIVA capsules (see **What the nonmedicinal ingredients are**).
- SUSTIVA should not be taken with some other medicines that are listed in this pamphlet (see the section entitled “**Drugs that may interact with SUSTIVA**”).

#### What the medicinal ingredient is:

Efavirenz

#### What the nonmedicinal ingredients are:

- SUSTIVA capsules also contain the following other inactive ingredients: sodium starch glycolate, lactose monohydrate, sodium lauryl sulfate and magnesium stearate. The capsule shell contains as excipients: gelatin, sodium lauryl sulfate, titanium dioxide and/or yellow iron oxide and may contain silicon dioxide. The capsules are printed with ink containing carmine, FD&C Blue No. 2 and titanium dioxide.
- 

#### What dosage forms it comes in:

- Each SUSTIVA capsule contains either 50 mg or 200 mg of efavirenz.

### WARNINGS AND PRECAUTIONS

#### BEFORE TAKING SUSTIVA

##### What should I tell my doctor before I start SUSTIVA?

- Inform your doctor about any past or present medical problems, including liver disease, hepatitis, allergies, severe kidney failure, seizures or mental illness.
- Inform your doctor if you have ever had a previous life-threatening skin reaction (e.g., Stevens-Johnson syndrome).
- Inform your doctor about any medications (prescription and nonprescription), herbal products, vitamins, nutritional supplements that you are currently taking or are planning to take.
- Also inform your doctor about any recreational (street, illicit) drugs that you are currently taking or are planning to take. The effect of combining

recreational (street, illicit) drugs or alcohol with SUSTIVA has not been studied. Because they may interact with each other, speak with your doctor or other healthcare provider before you combine these drugs.

- Inform your doctor if you have or have had a heart rhythm disorder such as an irregular heartbeat, prolongation of the QT interval or any risk factors for Torsade de Pointes (dangerous fluttering of the heart).

What should I consider concerning contraception, pregnancy, or breastfeeding?

- Tell your doctor if you are pregnant or planning to become pregnant. Birth defects have been reported in the offsprings of animals and women treated with SUSTIVA during pregnancy. It is not known whether SUSTIVA caused these defects. Women should not become pregnant while taking SUSTIVA and for 12 weeks after stopping it. If you are pregnant, you should take SUSTIVA only if you and your doctor decide that the possible benefit to you is greater than the possible risk to your foetus. If you take SUSTIVA while you are pregnant, talk to your doctor about how you can be included in the antiretroviral pregnancy registry.
- It is currently recommended that women living with HIV should not breastfeed because HIV infection can be passed on the baby in breast milk. If you are breastfeeding or thinking about breastfeeding you **should discuss it with** your doctor **as soon as possible**.
- A reliable form of barrier contraception must always be used even if you or your partner are using other methods of contraception such as the pill or other hormonal therapy (e.g., implants, injections). SUSTIVA may remain in your blood for a time after therapy is stopped. Therefore, you should continue use of a reliable form of contraception for 12 weeks after stopping treatment with SUSTIVA.

Can children take SUSTIVA?

- SUSTIVA has not been studied in children below 3 years of age.
- SUSTIVA can be taken by children 3 years or older and who are able to swallow capsules. Your child's doctor will determine the right dose based on your child's weight.

Do not drive or operate machinery until you have determined your response to SUSTIVA, as this may make you sleepy or dizzy.

SUSTIVA can cause abnormal blood test results. Your doctor may perform blood tests and will interpret the results.

To find out how to take SUSTIVA please read carefully the following section “**WHILE TAKING SUSTIVA**”.

**INTERACTIONS WITH THIS MEDICATION**

**Drugs that may interact with SUSTIVA**

SUSTIVA may affect the dosing of other medications including ones for treating HIV infection. For this reason it is very important to:

- Let all healthcare providers know that you are taking SUSTIVA.
- Inform your doctor and pharmacist about all medications that you are currently taking including those obtained over-the-counter without a prescription and complementary medications (vitamins, nutritional supplements, etc.) and herbal products, particularly St. John’s Wort.
- Consult your doctor or pharmacist before you start any new medication.
- Consult your doctor or pharmacist before you stop any medications that you are currently taking.

Bring all your medications when you see your doctor. Or make a list of their names, how much you take, and how often you take them. This will give your doctor a complete picture of the medications you are taking. Then he or she can decide the best approach for your situation.

**You must not take the following medications if you are taking SUSTIVA. Taking these medications with SUSTIVA could create the potential for serious and/or life-threatening side effects:**

- CISAPRIDE\*\***
- MIDAZOLAM**
- TRIAZOLAM (e.g, HALCION\*)**
- ERGOT MEDICATIONS (e.g, CAFERGOT\*)**
- PIMOZIDE (e.g, ORAP\*)**
- ZEPATIER**

**ST. JOHN’S WORT (*Hypericum perforatum*)**  
You must not take products containing St. John’s wort (*Hypericum perforatum*) with SUSTIVA as this may stop SUSTIVA from working properly and may lead to resistance to SUSTIVA or resistance to the class of non-nucleoside reverse transcriptase inhibitors (NNRTIs).

\* CISAPRIDE is not marketed in Canada

**The following drugs may interact with SUSTIVA and your doctor will determine whether they can be used or not or may make dosage changes for**

**SUSTIVA or the other product, or substitute other products, as indicated below:**

- SUSTIVA may be taken with many of the medications commonly used in people with HIV infection. These include the protease inhibitors, such as nelfinavir (Viracept<sup>®</sup>) and indinavir (Crixivan<sup>®</sup>), and nucleoside analogue reverse transcriptase inhibitors (NRTIs).
- **Use of SUSTIVA with saquinavir (Invirase<sup>®</sup>) is not recommended if you are taking saquinavir as your only protease inhibitor.**
- **VORICONAZOLE (VFEND<sup>®</sup>)** should not be taken at standard doses with SUSTIVA since it may lose its effect or increase the chance of side effects from SUSTIVA. Some doses of voriconazole can be taken at the same time as a lower dose of SUSTIVA, but your doctor will decide if this is appropriate.
- Tegretol<sup>®</sup> (carbamazepine), Sporanox<sup>®</sup> (itraconazole), Posanol<sup>®</sup> (posaconazole) and REYATAZ (atazanavir sulfate), if this is not the first time you are receiving treatment for your HIV infection, may need to be replaced with another medicine when taken with SUSTIVA.
- SUSTIVA reduces the blood levels of clarithromycin (Biaxin<sup>®</sup>) and is associated with a higher incidence of rash; your doctor may consider giving you an alternative antibiotic.
- If you are taking SUSTIVA and REYATAZ (atazanavir sulfate), you should also be taking Norvir<sup>®</sup> (ritonavir).
- Antimalarials such as atovaquone/proguanil, when taken with SUSTIVA, may reduce the amount of atovaquone/proguanil in your blood which may reduce the anti-malarial activity of these medicines. Atovaquone/proguanil should not be taken with SUSTIVA; your doctor should consider alternatives to these antimalarial medicines.
- Drugs that may interact with SUSTIVA to affect the electrical activity of your heart which include but may not be limited to macrolide antibiotics (such as clarithromycin) and antimalarials (artemether/lumefantrine).
- Use of EPCLUSA (sofosbuvir/velpatasvir), VOSEVI (sofosbuvir/velpatasvir/voxilaprevir), and MAVIRET (glecaprevir/pibrentasvir) is not recommended when taking SUSTIVA.
- SUSTIVA may interact with etonogestrel contraceptive implants. This means your implant might not work and you could get pregnant. You should use a barrier birth control method like a condom. Talk to your healthcare professional for advice on additional birth control methods.

**The following medicine should not be taken with**

**SUSTIVA since it contains efavirenz, the active ingredient in SUSTIVA:**

- ATRIPLA (efavirenz, emtricitabine, tenofovir disoproxil fumarate) unless your doctor decides a dose adjustment is needed (e.g. with rifampin).
- Your doctor may need to adjust the dose of either SUSTIVA or the following medications when taken with SUSTIVA:
  - Crixivan<sup>®</sup> (indinavir)
  - Methadone
  - Zoloft<sup>®</sup> (sertraline)
  - Wellbutrin SR, Wellbutrin XL, or Zyban (bupropion)
  - Kaletra<sup>®</sup> (lopinavir/ritonavir) Lopinavir and ritonavir combination should not be taken once daily with SUSTIVA. Your doctor may suggest an alternate dosing regimen.
  - Celsentri<sup>®</sup> (maraviroc)
  - Mycobutin<sup>®</sup> (rifabutin)
  - The cholesterol-lowering medicines Lipitor<sup>®</sup> (atorvastatin), Pravachol<sup>®</sup> (pravastatin), and Zocor<sup>®</sup> (simvastatin)
  - Rifadin<sup>®</sup> (rifampin) or the rifampin-containing medicines Rofact<sup>®</sup> and Rifater<sup>®</sup>
  - Calcium channel blockers such as Cardizem<sup>®</sup> or Tiazac<sup>®</sup> (diltiazem), Covera HS, Isoptin SR or Tarka (verapamil), and others.
  - Immunosuppressants such as Neoral<sup>®</sup> (cyclosporin), Advagraf<sup>®</sup> or Prograf<sup>®</sup> (tacrolimus), Rapamune<sup>®</sup> or Torisel<sup>®</sup> (sirolimus)
  - Hepatitis C antiviral agents
  - Antimalarials such as Coartem<sup>®</sup> ♦ and Riamet<sup>®</sup> ♦ (artemether/lumefantrine) *Anthelmintics* such as Biltricide<sup>®</sup> (praziquantel). Your doctor may have to consider increasing the dose of praziquantel.

♦ Not marketed in Canada

- The effect of combining alcohol or recreational (street, illicit) drugs with SUSTIVA has not been studied. Because they may interact with each other, speak with your doctor or other healthcare provider before you combine these drugs.

**PROPER USE OF THIS MEDICATION**

**WHILE TAKING SUSTIVA**

Usual Dose

- The dose of SUSTIVA for adults and children weighing more than 40 kg (88 lbs) is 600 mg once-a-day (three 200 mg capsules taken together).

- The dose for children weighing 40 kg or less is determined by the weight of the child and is taken once daily. SUSTIVA capsules should only be administered to children if they have the ability to reliably swallow capsules.
- You should take SUSTIVA on an empty stomach, preferably at bedtime. Taking SUSTIVA with food increases the level of efavirenz in the blood and may increase the possibility of side effects.
- Your doctor or pharmacist will give you instructions for proper dosage.

**What should I remember to do or avoid while taking SUSTIVA?**

- Swallow SUSTIVA with water.
- Do not chew the capsules.
- Taking SUSTIVA at bedtime may improve the tolerability of the nervous system side effects.
- It is important to take SUSTIVA as your doctor prescribes. Do not change the dose on your own.
- SUSTIVA should not be used alone to treat HIV. SUSTIVA should always be taken with other anti-HIV medications in order to prevent the virus from becoming resistant to your drug treatment.
- You should not stop taking SUSTIVA without first consulting with your doctor.
- If you are unsure of what to do or need help in planning the best times to take your medications, ask your doctor or other healthcare provider.
- If you think it would be useful, ask a friend or family member to remind you to take your medications.
- When your SUSTIVA supply starts to run low, get more from your doctor or pharmacist. This is very important because the amount of virus may start to increase if SUSTIVA is stopped for even a short time. The virus may then become harder to treat.
- Remember, SUSTIVA has been prescribed just for you. Never give your medications to others to try.
- Do not use your current supply of SUSTIVA after the end of the month and year shown by the “expiry date” on the bottle.

Overdose:

In case of drug overdosage, contact a healthcare practitioner (e.g., doctor), hospital emergency department or regional poison control centre, even if there are no symptoms.

Missed Dose:

- If you forget to take SUSTIVA, **do not double**

**your next dose.** Take the missed dose as soon as possible, and then carry on with your regular dosing schedule.

- Try not to miss a dose. With anti-HIV medications, missing doses or not taking them properly may allow the amount of HIV in your body to increase. HIV may then become resistant. This means that the virus changes or mutates causing a medication to lose its effect.

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

- SUSTIVA, like all medications, affects different people in different ways. Any medication may have unintended or unwanted effects, so-called side effects. Some people may develop side effects, others may not.
- The most notable side effects of SUSTIVA are rash and nervous system symptoms that include dizziness, insomnia (difficulty falling asleep), drowsiness, reduced ability to concentrate, and abnormal dreaming. These side effects are generally mild to moderate and tend to disappear after you have taken SUSTIVA for a few weeks. Decreasing the dose does not seem to help and is not recommended.
- Some of these side effects such as dizziness will likely be less noticeable if you take SUSTIVA before going to bed. Be sure to tell your doctor if any of these side effects continue or if they bother you.
- A small number of patients have had severe depression, strange thoughts, or angry behavior. Some patients have had thoughts of suicide and a few patients have actually committed suicide. These problems tend to occur more often in patients with a history of mental illness. You should contact your doctor immediately if you think you are having these symptoms, so your doctor can decide whether you should continue to take SUSTIVA.
- Dizziness, trouble concentrating, and drowsiness have been reported with SUSTIVA. If you notice any of these symptoms, you should avoid potentially hazardous tasks such as driving or operating machinery.
- You should consult your doctor if you have a rash since some rashes may be serious. However, most cases of rash disappear without any change in your treatment.
- Rash seems to be more common in children than in adults treated with SUSTIVA.
- Some patients taking SUSTIVA have experienced serious liver problems including liver failure

resulting in transplantation or death. Most of these serious side effects occurred in patients with a chronic liver disease such as a hepatitis infection, but there have also been a few reports in patients without any existing liver disease.

- Changes in your immune system (Immune Reconstitution Inflammatory Syndrome) can happen when you start taking medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Autoimmune disease (when the immune system attacks healthy body tissue) may also occur after you start taking medicines for HIV infection. Examples of this include the following conditions: Grave's disease, Guillain-Barre syndrome, polymyositis or autoimmune hepatitis ]. Autoimmune disorders may occur many months after the start of treatment. Look for any symptoms such as:
  - fever, redness, rash or swelling
  - fatigue
  - joint or muscle pain
  - numbness or weakness beginning in the hands and feet and moving up towards the trunk of the body
  - palpitations (chest pain) or rapid heart rate
  - bulging eyes, light sensitivity, or vision changes
  - yellowing of the skin
  - difficulty talking, chewing, or swallowing

If you notice any of these symptoms, tell your doctor immediately.

**Other side effects**

- Other common side effects that have been reported include tiredness, nausea, diarrhea and headache. These may be from SUSTIVA or from other medications that you are taking.
- Tell your doctor or other healthcare provider if you notice these or any other side effects not mentioned in the pamphlet that continue or if they bother you.

Remember do not stop taking SUSTIVA without speaking to your doctor first. He or she may be able to help you manage these side effects without stopping your anti-HIV medications.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM				
Effect/Symptoms		Talk with your doctor or pharmacist immediately		Stop taking drug and seek medical advice immediately
		Only if severe	In all cases	
Common	<b>Serious psychiatric events</b> <b>Symptoms:</b> <ul style="list-style-type: none"> <li>• Severe depression</li> <li>• Thoughts of suicide</li> <li>• Strange thoughts</li> <li>• Angry behavior.</li> <li>• Catatonia (unable to move or speak normally)</li> </ul>		✓	
	<b>Severe skin rash</b> <b>Symptoms:</b> <ul style="list-style-type: none"> <li>• Blisters or peeling of the skin</li> <li>• Blisters or peeling of the mouth, lips and throat</li> <li>• Fever and general ill feeling.</li> </ul>			✓
	<b>Liver failure</b> <b>Symptoms:</b> <ul style="list-style-type: none"> <li>• Jaundice (skin or the white part of eyes turn yellow)</li> <li>• Urine turns dark</li> <li>• Bowel movements (stools) turn light in color</li> <li>• Loss of appetite for several days or longer</li> <li>• Feeling sick to your stomach (nausea)</li> <li>• Lower stomach pain.</li> </ul>		✓	

*This is not a complete list of side effects. If you have any unexpected effects while taking SUSTIVA, contact your doctor or pharmacist.*

**HOW TO STORE IT**

SUSTIVA should be stored at room temperature (25°C; although a range of 15 – 30°C is permitted).

**As with all medications, SUSTIVA should be kept out of the reach and sight of children.**

**REPORTING SUSPECTED SIDE EFFECTS**

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect)
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
  - Fax toll-free to 1-866-678-6789, or
  - Mail to: Canada Vigilance Program  
Health Canada  
Postal Locator 0701C  
Ottawa, ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reactions reporting guidelines are available on the MedEffect™ Canada Web site at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect).

*Note: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.*

**MORE INFORMATION**

This document plus the full product monograph, prepared for health professionals can be obtained by contacting the sponsor, Bristol-Myers Squibb Canada, at:  
1-866-463-6267.

This leaflet was prepared by Bristol-Myers Squibb Canada.

Last revised: 16 August 2023

SUSTIVA® is a registered trademark of Bristol-Myers Squibb Pharma Company used under license by Bristol-Myers Squibb Canada.

REYATAZ® is a registered trademark of Bristol-Myers Squibb Holdings Ireland used under license by Bristol-Myers Squibb Canada.

PRAVACHOL® is a registered trademark of Bristol-Myers Squibb Company used under license by Bristol-Myers Squibb Canada.

\*Other brands listed are trademarks of their respective owners.