

BREYANZI®

LISOCABTAGENE MARALEUCEL (LISO-CEL)

PATIENT CARD

Have this card with you at all times. Show it to any doctor/healthcare provider who sees you, including in an emergency.

Tell any healthcare provider who sees you that you are being treated with liso-cel.

For at least 2 weeks after receiving liso-cel, you should plan to stay close (within approximately a 2 hour drive) to the location where you received treatment. Your doctor may recommend that you stay close by for a longer period.

For at least 4 weeks after receiving liso-cel, you should not drive, use machines, or take part in activities that need you to be alert. Based on your individual needs, your doctor may advise you to wait longer before driving.

I have been treated with liso-cel

Important Contact Information

My name: _____

Name of liso-cel treating physician: _____

Hospital phone number: _____

Hospital phone number outside office hours: _____

Hospital name: _____

Date of liso-cel infusion (dd-mm-yyyy): _____

Batch number: _____

Information for the Healthcare Provider

This patient has received liso-cel chimeric antigen receptor (CAR)-positive T-cell therapy.

Following treatment with liso-cel, cytokine release syndrome (CRS) and/or neurologic toxicities, including immune effector cell-associated neurotoxicity syndrome (ICANS), may occur, which may be fatal or life-threatening. Cytokine release syndrome (CRS) may involve any organ system.

Contact patient's liso-cel treating physician immediately for further information.

Please see liso-cel's Summary of Product Characteristics and Package leaflet which can be found at www.geneesmiddeleninformatiebank.nl.

Information for Patients

Liso-cel may cause serious side effects in various parts of your body. These side effects can be life-threatening or even fatal and must be treated immediately.

Symptoms that seem mild can worsen quickly.

Symptoms can develop later, sometimes weeks to months after your infusion.

Call your liso-cel treating physician or go to the emergency room/department immediately if any of the following symptoms appear:

Neurologic adverse reactions

Cytokine release syndrome (CRS)

The following may be symptoms of ICANS:

- Confusion
- Being less alert (decreased consciousness)
- Difficulty speaking or slurred speech
- Shaking (tremor)
- Feeling anxious
- Feeling dizzy
- Headache
- Fever
- Chills or shaking
- Feeling tired
- Fast or uneven heartbeat
- Feeling light-headed and short of breath

Would you like more information about your medication?

Read the package insert or scan the QR code for online resources. If you have any questions about your medication or side effects, please contact your doctor or pharmacist.

